



Laboratoire d'Annecy-le-Vieux de Physique Théorique

website: <http://lappweb.in2p3.fr/lapth-2005/>

Nested Bethe ansatz for 'all' open spin chains with diagonal boundary conditions

S. Belliard¹ and E. Ragoucy²

Laboratoire de Physique Théorique LAPTh

UMR 5108 du CNRS, associée à l'Université de Savoie

BP 110, F-74941 Annecy-le-Vieux Cedex, France.

Abstract

We present in an unified and detailed way the nested Bethe ansatz for open spin chains based on $\mathcal{V}(gl(\mathbf{n}))$, $\mathcal{V}(gl(\mathbf{m}|\mathbf{n}))$, $\hat{\mathcal{U}}_q(gl(\mathbf{n}))$ or $\hat{\mathcal{U}}_q(gl(\mathbf{m}|\mathbf{n}))$ (super)algebras, with arbitrary representations (i.e. 'spins') on each site of the chain and diagonal boundary matrices $(K^+(u), K^-(u))$. The nested Bethe ansatz applies for a general $K^-(u)$, but a particular form of the $K^+(u)$ matrix.

The construction extends and unifies the results already obtained for open spin chains based on fundamental representation and for some particular super-spin chains. We give the eigenvalues, Bethe equations and the explicit form of the Bethe vectors for the corresponding models. The Bethe vectors are also expressed using a trace formula.

MSC: 81R50, 17B37 — PACS: 02.20.Uw, 03.65.Fd, 75.10.Pq

LAPTh-1309/09

arXiv:0902.0321 [math-ph]

February 2009

¹samuel.belliard@lapp.in2p3.fr

²eric.ragoucy@lapp.in2p3.fr

1 Introduction

The systematic studies of the open spins chains using R matrices formalism start with the seminal papers of Cherednik [1] and Sklyanin [2], who generalized to these models the QISM approach developed by the Leningrad school. They introduced the reflection algebra as the fundamental ingredient to construct the Abelian Bethe subalgebra and ensure integrability of the model. This algebra is a subalgebra of the FRT algebra introduced by Leningrad group for the periodic spin chains (for a review, see e.g. [3] and references therein). The boundary conditions are encoded in two matrices, $K^-(u)$ solution of the reflection equation, see equation (3.8) below, and $K^+(u)$ solution of the dual equation, see equation (3.14). With these matrices and the standard closed spin chain monodromy matrix, one can construct a transfer matrix that belongs to the Bethe subalgebra. The existence of this subalgebra lead to the integrability of the model when the expansion of the transfer matrix as a series provides a sufficient number of operators in involution. In the following we consider that this number is sufficient.

After proving integrability of the model, the next step is to find the eigenvalues and eigenvectors of this Bethe subalgebra. It depends on the choice of the boundary matrices. Focusing on diagonalizable boundary matrices, two main cases can be distinguished: $K^+(u)$ and $K^-(v)$ are diagonalizable in the same basis; or not.

Very few is known in the latter case, apart from two recent approaches developed for XXZ spin chain and that does not rely on (nested) algebraic Bethe ansatz [4–6]: in [7], the reflection equation is replaced by a deformed Onsager algebra (which may be another presentation of the reflection algebra); and in [8], eigenvalues are computed using generalized TQ relations when $K^-(u)$ and $K^+(u)$ obey some relations, or when the deformation parameter is root of unity.

The first case can be divided into two sub-families, depending whether (i) the diagonalization matrix is a constant or (ii) depends on the spectral parameter. Again, in the case (ii), only some results are known from the gauge transformation construction of [9–11], that allows to relate non-diagonal solutions to diagonal ones via a Face-Vertex correspondence. The case (i) is the one studied by analytical Bethe ansatz [12] and corresponds to diagonal matrices

$$K^-(u) = \text{diag}(\underbrace{k^-(u), \dots, k^-(u)}_{a_-}, \underbrace{\bar{k}^-(u), \dots, \bar{k}^-(u)}_{m+n-a_-}), \quad (1.1)$$

$$K^+(u) = \text{diag}(\underbrace{k^+(u), \dots, k^+(u)}_{a_+}, \underbrace{\bar{k}^+(u), \dots, \bar{k}^+(u)}_{m+n-a_+}). \quad (1.2)$$

Indeed, using this ansatz, eigenvalues of the transfer matrix can be computed for all (open or closed) chains based on $gl(\mathbf{n})$ and $gl(\mathbf{m}|\mathbf{n})$ (super)algebras and their deformation, and with arbitrary representations on each sites [13–15]. It is in general believed that a Nested (algebraic) Bethe Ansatz (NBA) can provide the eigenvectors of the corresponding models. This was shown in a unified way in [16] for closed spin chains.

We present here the open spin chains case. We will show that the standard NBA approach does not work in the general case. Keeping a general diagonal solution for $K^-(u)$, one needs to take $a_+ = 0$ or $a_+ = 1$ to perform a complete nested algebraic Bethe ansatz. In this case, the couple $(K^-(u), K^+(u))$ will be called a **NABA couple**. When one studies an open spin chain possessing a couple of diagonal matrices $(K^-(u), K^+(u))$ that is not of type NABA, one can start the first step of NBA approach, but then needs to switch (and end) the calculation with an analytical Bethe ansatz, as it has been done in e.g. [10, 11, 17].

In the present paper, we focus on NABA couples. Performing NBA, we compute the Bethe ansatz equations, the eigenvalues and the eigenvectors of the corresponding transfer matrix and show where the constraint $a_+ = 0$ or $a_+ = 1$ is needed in the calculation. Our presentation consider **universal transfer matrices** in the sense that the calculation applies to transfer matrices based on $gl(\mathbf{n})$ and $gl(\mathbf{m}|\mathbf{n})$ algebras and their deformation, with any finite dimensional irreducible representations of the monodromy matrix. In particular, it encompasses the preview results obtained for fundamental representations [10, 17, 18].

In addition to the derivation of the Bethe ansatz equations and transfer matrix spectrum, our main result is the explicit construction of the Bethe vectors. This is reflected in e.g. the trace formula (see theorem 7.1 at the end of the paper).

The plan of the paper is as follows. In section 2, we introduce the different notations and R -matrices we use in the paper. Then, in section 3, we present, using the FRT [19] formalism, the algebras concerned with our approach. They are generalizations of loop algebras (quantum algebras or Yangians, and their graded versions) and noted $\mathcal{A}_{\mathbf{m}|\mathbf{n}}$. They contain as subalgebras the reflection algebras, noted $\mathfrak{D}_{\mathbf{m}|\mathbf{n}}$. We also construct mappings

$$\mathfrak{D}_{\mathbf{m}|\mathbf{n}} \rightarrow \mathfrak{D}_{\mathbf{m}-1|\mathbf{n}} \rightarrow \cdots \rightarrow \mathfrak{D}_{1|1} \text{ or } \mathfrak{D}_2$$

that are needed for the nesting. In section 4, we present the finite dimensional irreducible representations of $\mathcal{A}_{\mathbf{m}|\mathbf{n}}$ and we compute the form of $T^{-1}(u)$. We also construct the representations of $\mathfrak{D}_{\mathbf{m}|\mathbf{n}}$ from the $\mathcal{A}_{\mathbf{m}|\mathbf{n}}$ ones. In section 5, as a warm up, we recall the algebraic Bethe ansatz, which deals with spin chains based on $gl(2)$, $gl(1|1)$ algebras and their quantum deformations. Then, in section 6, we perform the nested Bethe ansatz in a very detailed and pedestrian way and up to the end. Finally, in section 7, we study the Bethe vectors that have been constructed in the previous section, showing connection with a **trace formula**. As a conclusion, we discuss our results and present some possible applications or extensions of our work.

2 Notations

2.1 Graded auxiliary spaces

We use the so-called auxiliary space framework. In this formalism, one deals with multiple tensor product of vectorial spaces $\mathcal{V} \otimes \cdots \otimes \mathcal{V}$, and operators (defining an algebra $\mathcal{A}$) therein. For any matrix valued operator, $A := \sum_{ij} E_{ij} \otimes a_{ij} \in \text{End}(\mathcal{V}) \otimes \mathcal{A}$, we set

$$A_k := \sum_{ij} \mathbb{I}^{\otimes(k-1)} \otimes E_{ij} \otimes \mathbb{I}^{\otimes(m-k)} \otimes a_{ij} \in \text{End}(\mathcal{V}^{\otimes m}) \otimes \mathcal{A}, \quad 1 \leq k \leq m, \quad (2.1)$$

where E_{ij} are elementary matrices, with 1 at position (i, j) and 0 elsewhere. The notation is valid for complex matrices, taking $\mathcal{A} := \mathbb{C}$ and using the isomorphism $\text{End}(\mathcal{V}) \otimes \mathbb{C} \sim \text{End}(\mathcal{V})$.

We will work on $\mathbb{Z}_2$ -graded spaces $\mathbb{C}^{\mathbf{m}|\mathbf{n}}$. The elementary $\mathbb{C}^{\mathbf{m}|\mathbf{n}}$ column vectors e_i (with 1 at position i and 0 elsewhere) and elementary $\text{End}(\mathbb{C}^{\mathbf{m}|\mathbf{n}})$ matrices E_{ij} have grade:

$$[e_i] = [i] \quad \text{and} \quad [E_{ij}] = [i] + [j]. \quad (2.2)$$

This grading is also extended to the superalgebras we deal with, see section 3.1 below. The tensor product is graded accordingly:

$$(a_{ij} \otimes a_{kl})(a_{pq} \otimes a_{rs}) = (-1)^{([k]+[l])([p]+[q])}(a_{ij}a_{pq} \otimes a_{kl}a_{rs}). \quad (2.3)$$

The transposition $(.)^t$ and trace $str(.)$ operators are also graded:

$$A^t = \sum_{i,j=1}^{m+n} (-1)^{[j]+[j][i]} E_{ji} \otimes a_{ij}, \quad str A = \sum_{i=1}^{m+n} (-1)^{[i]} a_{ii} \quad \text{for} \quad A = \sum_{i,j=1}^{m+n} E_{ij} \otimes a_{ij}. \quad (2.4)$$

To simplify the presentation we work with the *distinguished* $\mathbb{Z}_2$ -grade defined by:

$$[i] = \begin{cases} 0, & 1 \leq i \leq m, \\ 1, & m+1 \leq i \leq m+n. \end{cases} \quad (2.5)$$

Simplification in the expressions follows from the rule $[i][j] = [i]$ when $i \leq j$, which is valid only for the distinguished grade. The non graded case is recovered setting $n = 0$ and $[k] = 0$.

2.2 Spectral parameters transformations

For spectral parameter u we use the following notations:

$$\begin{aligned} \iota(u) &= \begin{cases} -u & \text{for } \mathcal{Y}(\mathbf{m}|\mathbf{n}) \\ \frac{1}{u} & \text{for } \widehat{\mathcal{U}}_q(\mathbf{m}|\mathbf{n}) \end{cases} ; \quad \tilde{u} = \begin{cases} u - \frac{(m-n)\hbar}{2} & \text{for } \mathcal{Y}(\mathbf{m}|\mathbf{n}) \\ u q^{\frac{(m-n)}{2}} & \text{for } \widehat{\mathcal{U}}_q(\mathbf{m}|\mathbf{n}) \end{cases} ; \\ u^{(k)} &= \begin{cases} u + \frac{\hbar}{2}(-1)^{[k]} & \text{for } \mathcal{Y}(\mathbf{m}|\mathbf{n}) \\ u q^{-\frac{1}{2}+2[k]} & \text{for } \widehat{\mathcal{U}}_q(\mathbf{m}|\mathbf{n}) \end{cases} ; \quad u^{\overline{(k)}} = \begin{cases} u - \frac{\hbar}{2}(-1)^{[k]} & \text{for } \mathcal{Y}(\mathbf{m}|\mathbf{n}) \\ u q^{\frac{1}{2}-[k]} & \text{for } \widehat{\mathcal{U}}_q(\mathbf{m}|\mathbf{n}) \end{cases} ; \\ u^{(k\dots l)} &= (\dots (u^{(k)})^{(k+1)} \dots)^{(l)}. \end{aligned}$$

2.3 R -matrices

In what follows, we will deal with different types of matrices $R \in End(\mathcal{V}) \otimes End(\mathcal{V})$, all obeying (graded) Yang-Baxter equation (written in auxiliary space $End(\mathcal{V}) \otimes End(\mathcal{V}) \otimes End(\mathcal{V})$):

$$R_{12}(u_1, u_2) R_{13}(u_1, u_3) R_{23}(u_2, u_3) = R_{23}(u_2, u_3) R_{13}(u_1, u_3) R_{12}(u_1, u_2). \quad (2.6)$$

The R -matrix satisfy unitarity relation,

$$R_{12}(u, v) R_{21}(v, u) = \zeta(u, v) \mathbb{I} \otimes \mathbb{I}, \quad (2.7)$$

and crossing unitarity (see (2.11) below for the definition of $\bar{R}$),

$$R_{12}^{t_1}(u, v) M_1 \bar{R}_{12}^{t_2}(\iota(\tilde{v}), \iota(\tilde{u})) M_1^{-1} = \bar{\zeta}(u, v) \mathbb{I} \otimes \mathbb{I}, \quad (2.8)$$

where $\zeta(u, v)$ and $\bar{\zeta}(u, v)$ are $\mathbb{C}$ -functions depending on the model under consideration, M is a $\mathbb{C}$ -valued matrix defined in appendix A for each model and t_a is the transposition in the auxiliary space a . All the R -matrices used here also obey the parity relation:

$$R_{12}(u, v)^{t_1 t_2} = R_{21}(u, v). \quad (2.9)$$

To each R -matrix, one associates an algebra $\mathcal{A}_{\mathbf{m}|\mathbf{n}}$ using the FRT formalism. Below, we focus on infinite dimensional associative algebras based on $gl(\mathbf{n})$ and $gl(\mathbf{m}|\mathbf{n})$ Lie (super)algebras and

their q -deformation. We note these algebras $\mathcal{A}_{\mathbf{n}} = Y(\mathbf{n})$ or $\widehat{\mathcal{U}}_q(\mathbf{n})$ and $\mathcal{A}_{\mathbf{m}|\mathbf{n}} = \mathcal{Y}(\mathbf{m}|\mathbf{n})$ or $\widehat{\mathcal{U}}_q(\mathbf{m}|\mathbf{n})$. We will write also $\mathcal{A}_{\mathbf{m}|0} = \mathcal{A}_{\mathbf{m}}$. We will encompass all R -matrices of these algebras writing:

$$R_{12}(u, v) = \mathbf{b}(u, v) \mathbb{I} \otimes \mathbb{I} + \sum_{i,j=1}^{\mathbf{m}+\mathbf{n}} \mathbf{w}_{ij}(u, v) E_{ij} \otimes E_{ji}. \quad (2.10)$$

All functions are defined in appendix B for each cases (one can refer to [16] for details and references). To define the reflection equation, we need another R -matrix:

$$\bar{R}_{12}(u, v) = R_{12}(u, \iota(v)) = \bar{\mathbf{b}}(u, v) \mathbb{I} \otimes \mathbb{I} + \sum_{i,j=1}^{\mathbf{m}+\mathbf{n}} \bar{\mathbf{w}}_{ij}(u, v) E_{ij} \otimes E_{ji} \quad (2.11)$$

From (2.6) we can deduce the relation between these two R matrices:

$$R_{12}(u_1, u_2) \bar{R}_{13}(u_1, u_3) \bar{R}_{23}(u_2, u_3) = \bar{R}_{23}(u_2, u_3) \bar{R}_{13}(u_1, u_3) R_{12}(u_1, u_2). \quad (2.12)$$

We will also use ‘reduced’ R -matrices $R^{(k)}(u)$, deduced from $R(u)$ by suppressing all the terms containing indices j with $j < k$:

$$\begin{aligned} R_{12}^{(k)}(u, v) &= (\mathbb{I}^{(k)} \otimes \mathbb{I}^{(k)}) R_{12}(u, v) (\mathbb{I}^{(k)} \otimes \mathbb{I}^{(k)}) \\ &= \mathbf{b}(u, v) \mathbb{I}^{(k)} \otimes \mathbb{I}^{(k)} + \sum_{i,j=k}^{\mathbf{m}+\mathbf{n}} \mathbf{w}_{ij}(u, v) E_{ij} \otimes E_{ji}, \end{aligned} \quad (2.13)$$

$$\text{where} \quad \mathbb{I}^{(k)} = \sum_{i=k}^{\mathbf{m}+\mathbf{n}} E_{ii}, \quad \forall k. \quad (2.14)$$

$R_{12}^{(k)}(u, v)$ corresponds to the R -matrix of³ $\mathcal{A}_{\mathbf{m}+1-k|\mathbf{n}}$. We will also use:

$$\begin{aligned} R_{12}^{(k,p)}(u, v) &= (\mathbb{I}^{(k)} \otimes \mathbb{I}^{(p)}) R_{12}(u, v) (\mathbb{I}^{(k)} \otimes \mathbb{I}^{(p)}) \\ &= \mathbf{b}(u, v) \mathbb{I}^{(k)} \otimes \mathbb{I}^{(p)} + \sum_{i,j=\max(k,p)}^{\mathbf{m}+\mathbf{n}} \mathbf{w}_{ij}(u, v) E_{ij} \otimes E_{ji}. \end{aligned} \quad (2.15)$$

Note that $R_{12}^{(k,k)}(u, v) = R_{12}^{(k)}(u, v)$. We define the ‘normalized reduced’ R -matrices:

$$\mathbb{R}_{12}^{(k,p)}(u, v) = \frac{1}{\mathbf{a}_p(u, v)} R_{12}^{(k,p)}(u, v) \quad \text{with} \quad \mathbb{R}_{12}^{(k,k)}(u, v) \mathbb{R}_{21}^{(k,k)}(v, u) = \mathbb{I}^{(k)} \otimes \mathbb{I}^{(k)}. \quad (2.16)$$

3 Algebraic structures

3.1 FRT formalism

The FRT (or RTT) relations [19–21] allow us to generate all the relations between the generators of the graded unital associative algebra $\mathcal{A}_{\mathbf{m}|\mathbf{n}}$. We gather the $\mathcal{A}_{\mathbf{m}|\mathbf{n}}$ generators into a $(\mathbf{m} + \mathbf{n}) \times$

³We will write, for a generic k , $\mathbf{m} - k|\mathbf{n}$, keeping in mind that one should write $0|\mathbf{n} - (k - \mathbf{m})$ when $k > \mathbf{m}$.

$(\mathfrak{m} + \mathfrak{n})$ matrix acting in an auxiliary space $\mathcal{V} = \mathbb{C}^{\mathfrak{m}|\mathfrak{n}}$ whose entries are formal series of a complex parameter u :

$$T(u) = \sum_{i,j=1}^{\mathfrak{m}+\mathfrak{n}} E_{ij} \otimes t_{ij}(u) \in \text{End}(\mathcal{V}) \otimes \mathcal{A}[[u, u^{-1}]]. \quad (3.1)$$

Since the auxiliary space $\text{End}(\mathbb{C}^{\mathfrak{m}|\mathfrak{n}})$ is interpreted as a representation of $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$, the $\mathbb{Z}_2$ -grading of $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ must correspond to the one defined on $\text{End}(\mathbb{C}^{\mathfrak{m}|\mathfrak{n}})$ matrices (see section 2). Hence, the generator $t_{ij}(u)$ has grade $[i] + [j]$, so that the monodromy matrix $T(u)$ is globally even. As for matrices, the tensor product of algebras will be graded, as well as between algebras and matrices,

$$(E_{ij} \otimes t_{ij}(u)) (E_{kl} \otimes t_{kl}(v)) = (-1)^{([i]+[j])([k]+[l])} E_{ij} E_{kl} \otimes t_{ij}(u) t_{kl}(v). \quad (3.2)$$

The ‘real’ generators $t_{ij}^{(n)}$ of $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ appear upon expansion of $t_{ij}(u)$ in u . For the (super) Yangians $\mathcal{Y}(\mathfrak{n})$ and $\mathcal{Y}(\mathfrak{m}|\mathfrak{n})$, $t_{ij}(u)$ is a series in u^{-1} :

$$t_{ij}(u) = \sum_{n=0}^{\infty} t_{ij}^{(n)} u^{-n} \quad \text{with} \quad t_{ij}^{(0)} = \delta_{ij}. \quad (3.3)$$

In the quantum affine (super)algebra [22,23] without central charge case, a complete description of the algebras requires the introduction of two matrices $L^{\pm}(u)$. However, in the context of evaluation representations it is sufficient to consider only $T(u) = L^{+}(u)$ to construct a transfer matrix. Indeed, in an evaluation representation, the choices $T(u) = L^{-}(u)$ or $T(u) = L^{+}(u) - L^{-}(u)$ lead to the same operator up to a multiplication function. Then, the RTT relations take the form:

$$R_{12}(u, v) T_1(u) T_2(v) = T_2(v) T_1(u) R_{12}(u, v). \quad (3.4)$$

$\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ has the following antimorphisms:

$$\begin{aligned} \text{Matrix inversion} \quad \text{inv} : \quad T(u) &\rightarrow T^{-1}(u) = \sum_{i,j=1}^{\mathfrak{m}+\mathfrak{n}} E_{ij} \otimes t'_{ij}(u), \\ \text{Matrix transposition} \quad t : \quad T(u) &\rightarrow T^t(u) = \sum_{i,j=1}^{\mathfrak{m}+\mathfrak{n}} (E_{ij})^t \otimes t_{ij}(u), \\ \text{Spectral parameter inversion} \quad \iota : \quad T(u) &\rightarrow T(\iota(u)). \end{aligned} \quad (3.5)$$

$\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ has a Hopf algebra structure, with coproduct

$$\Delta(T(u)) = T(u) \dot{\otimes} T(u) = \sum_{i,j,k=1}^{\mathfrak{m}+\mathfrak{n}} (-1)^{([k]+[i])([k]+[j])} E_{ij} \otimes t_{ik}(u) \otimes t_{kj}(u). \quad (3.6)$$

More generally, one defines recursively for $L \geq 2$, the algebra homomorphism

$$\Delta^{(L+1)} = (\text{id}^{\otimes(L-1)} \otimes \Delta) \circ \Delta^{(L)} \quad \text{with} \quad \Delta^{(2)} = \Delta \quad \text{and} \quad \Delta^{(1)} = \text{id}. \quad (3.7)$$

3.2 Reflection algebra and $K(u)$ matrices

The $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ algebra is enough to construct a transfer matrix leading to periodic spin chain models. In the context of open spin chains, one needs another algebra, the reflection algebra $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$, which turns out to be a subalgebra of $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$. Indeed, physically, one can interpret the FRT relation as encoding the interaction between the spins of the chain. Hence, it is the only relation needed to describe a periodic chain. On the other hand, in the case of open chain, the interaction with the boundaries has to be taken into account. Following the seminal paper of Sklyanin [2], we construct the reflection algebra and the dual reflection equation for the boundary scalar matrices $K^-(u)$ and $K^+(u)$. We first define the matrix $K^-(u)$ to be the solution of the reflection equation:

$$R_{12}(u_1, u_2) K_1^-(u_1) \bar{R}_{21}(u_1, u_2) K_2^-(u_2) = K_2^-(u_2) \bar{R}_{12}(u_1, u_2) K_1^-(u_1) R_{21}(u_1, u_2). \quad (3.8)$$

Depending on the type of R -matrix one considers, solutions to the reflection equation have been classified: see [24] for the Yangian and super-Yangian cases; in the other cases, partial classifications have been obtained in e.g. [17, 25]. In all cases, diagonal solutions of the reflection equations are known. They take the form (up to normalisation),

$$K^-(u) = \begin{cases} \text{diag}(\underbrace{u - c_-, \dots, u - c_-}_a, \underbrace{-u - c_-, \dots, -u - c_-}_{\mathfrak{m}+\mathfrak{n}-a}) & \text{for } \mathcal{Y}(\mathfrak{m}|\mathfrak{n}), \\ \text{diag}(\underbrace{u^2 - c_-^2, \dots, u^2 - c_-^2}_a, \underbrace{u^{-2} - c_-^2, \dots, u^{-2} - c_-^2}_{\mathfrak{m}+\mathfrak{n}-a}) & \text{for } \hat{\mathcal{U}}_q(\mathfrak{m}|\mathfrak{n}), \end{cases} \quad (3.9)$$

where c_- is a free complex parameter and a is an integer such that $0 \leq a \leq \mathfrak{m} + \mathfrak{n}$. From this $K^-(u)$ matrix and the monodromy matrix $T(u)$ for closed spin chains, we can construct the monodromy matrix of open spin chain:

$$D(u) = T(u) K^-(u) T^{-1}(\boldsymbol{\iota}(u)) = \sum_{i,j=1}^{\mathfrak{m}+\mathfrak{n}} E_{ij} \otimes d_{ij}(u), \quad (3.10)$$

$$d_{ij}(u) = \sum_{a=1}^{\mathfrak{m}+\mathfrak{n}} (-1)^{([i]+[a])([a]+[j])} \kappa_a(u) t_{ia}(u) t'_{aj}(\boldsymbol{\iota}(u)). \quad (3.11)$$

From (3.4) and (3.8), we can prove that $D(u)$ also satisfies the reflection equation:

$$R_{12}(u_1, u_2) D_1(u_1) \bar{R}_{21}(u_1, u_2) D_2(u_2) = D_2(u_2) \bar{R}_{12}(u_1, u_2) D_1(u_1) R_{21}(u_1, u_2). \quad (3.12)$$

This relation defines the reflection algebra $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$. The algebra $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ is a left coideal [26] of the algebra $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ with coproduct:

$$\Delta D_{[2]}(u) = T_{[1]}(u) D_{[2]}(u) T_{[1]}^{-1}(\boldsymbol{\iota}(u)) \in \text{End}(\mathcal{V}) \otimes \mathcal{A}_{\mathfrak{m}|\mathfrak{n}} \otimes \mathfrak{D}_{\mathfrak{m}|\mathfrak{n}} \quad (3.13)$$

where $[i]$ labels the two copies of $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$. This expression allows to increase the number of sites for an open spin chain in the same way one does for periodic ones: one acts on the monodromy matrix with the coproduct and then represents the new copy of algebra on the new ‘site’.

We also need a dual equation to construct transfer matrices in involution:

$$\begin{aligned} R_{12}(u_2, u_1)(K_1^+(u_1))^{t_1} M_1^{-1} \bar{R}_{21}(\boldsymbol{\iota}(\tilde{u}_1), \boldsymbol{\iota}(\tilde{u}_2)) M_1(K_2^+(u_2))^{t_2} = \\ (K_2^+(u_2))^{t_2} M_1 \bar{R}_{12}(\boldsymbol{\iota}(\tilde{u}_1), \boldsymbol{\iota}(\tilde{u}_2)) M_1^{-1} (K_1^+(u_1))^{t_1} R_{21}(u_2, u_1). \end{aligned} \quad (3.14)$$

where M is given in appendix A. From the property

$$R_{12}(u_1, u_2) M_1 M_2 = M_1 M_2 R_{12}(u_1, u_2), \quad (3.15)$$

one can construct solutions to the dual reflection equation using $K^-(u)$ solutions:

$$(K^+(u))^t = M K^-(\boldsymbol{\iota}(\tilde{u})). \quad (3.16)$$

With $D(u)$ and $K^+(u)$ we construct the transfer matrix:

$$d(u) = \text{str}(K^+(u)D(u)). \quad (3.17)$$

The reflection equation and its dual form ensure the commutation relation $[d(u), d(v)] = 0$. Thus, $d(u)$ generates (via an expansion in u) a set of L (the number of sites) independent integrals of motion (or charges) in involution which ensure integrability of the model.

3.2.1 Commutation relations of $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$

Projecting (3.12) on the $E_{ij} \otimes E_{kl}$ basis we get the commutation relations for $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$:

$$\begin{aligned} [d_{ij}(u), d_{kl}(v)] &= -\delta_{kj} \sum_{a=1}^{m+n} \frac{\bar{\mathfrak{w}}_{ka}(u, v)}{\bar{\mathfrak{b}}(u, v)} (-1)^{([i]+[l])([a]+[k])} d_{ia}(u) d_{al}(v) \\ &+ \delta_{il} \sum_{a=1}^{m+n} \frac{\bar{\mathfrak{w}}_{ia}(u, v)}{\bar{\mathfrak{b}}(u, v)} (-1)^{[l]+[a][k]+[a][j]+[k][j]} d_{ka}(v) d_{aj}(u) \\ &- \delta_{ij} \frac{\mathfrak{w}_{ik}(u, v)}{\mathfrak{b}(u, v)} \sum_{a=1}^{m+n} \frac{\bar{\mathfrak{w}}_{ia}(u, v)}{\bar{\mathfrak{b}}(u, v)} (-1)^{([a]+[k])([k]+[l])} d_{ka}(u) d_{al}(v) \\ &+ \delta_{ij} \frac{\mathfrak{w}_{il}(u, v)}{\mathfrak{b}(u, v)} \sum_{a=1}^{m+n} \frac{\bar{\mathfrak{w}}_{ia}(u, v)}{\bar{\mathfrak{b}}(u, v)} (-1)^{([a]+[l])([k]+[l])} d_{ka}(v) d_{al}(u) \\ &- \frac{\mathfrak{w}_{ik}(u, v)}{\mathfrak{b}(u, v)} (-1)^{([i]+[k])([k]+[l])} d_{kj}(u) d_{il}(v) + \frac{\mathfrak{w}_{jl}(u, v)}{\mathfrak{b}(u, v)} (-1)^{([i]+[l])([k]+[l])} d_{kj}(v) d_{il}(u) \end{aligned} \quad (3.18)$$

where $[x, y] = xy - (-1)^{[x][y]}yx$ is the graded commutator.

3.3 Embeddings of $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ algebras

The algebraic cornerstone for the nested Bethe ansatz is a recursion relation on the $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ algebraic structure. In this section we present a coset construction for $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ algebras (see theorem 3.1), that extends to the coideal property (see lemma 3.2 and theorem 3.3).

Theorem 3.1 For $k = 1, 2, \dots, \mathbf{m} + \mathbf{n} - 1$, let $F^{(k)}$ be a linear combination of elements $d_{i_1 j_1}(u_1) \dots d_{i_l j_l}(u_l)$ with all indices $i_p, j_p > k - 1$, and let $\mathcal{I}_k$ be the left ideal generated by $d_{ij}(u)$ for $i > j$ and $j < k$. Then, we have the following properties:

$$d_{ij}(u) F^{(k)} \equiv 0 \pmod{\mathcal{I}_k}, \quad \text{for } i > j \text{ and } j < k, \quad (3.19)$$

$$[d_{ii}(u), F^{(k)}] \equiv 0 \pmod{\mathcal{I}_k}, \quad \text{for } i < k. \quad (3.20)$$

Using the functions ψ_j given in (B.2), we introduce the generators:

$$\widehat{D}^{(k)}(u) = \sum_{i,j=k}^{\mathbf{m}+\mathbf{n}} E_{ij} \otimes d_{ij}^{(k)}(u), \quad (3.21)$$

$$d_{ij}^{(k)}(u) = d_{ij}(u^{(1\dots k-1)}) - \delta_{ij} \sum_{a=1}^{k-1} q^{2(k-1-a)-4 \sum_{l=a+1}^{k-1} [l]} \psi_a(u^{(a)}) d_{aa}(u^{(1\dots k-1)}). \quad (3.22)$$

They satisfy in $\mathfrak{D}_{\mathbf{m}|\mathbf{n}}/\mathcal{I}_k$ the reflection equation for $\mathfrak{D}_{\mathbf{m}-k+1|\mathbf{n}}$:

$$R_{12}^{(k)}(u_1, u_2) \widehat{D}_1^{(k)}(u_1) \bar{R}_{21}^{(k)}(u_1, u_2) \widehat{D}_2^{(k)}(u_2) \equiv \widehat{D}_2^{(k)}(u_2) \bar{R}_{12}^{(k)}(u_1, u_2) \widehat{D}_1^{(k)}(u_1) R_{21}^{(k)}(u_1, u_2) \pmod{\mathcal{I}_k}. \quad (3.23)$$

Proof: We first prove relation (3.20) for $k = 2$, the case $k = 1$ being trivially satisfied. A direct calculation from the commutation relations (3.18) of $\mathfrak{D}_{\mathbf{m}|\mathbf{n}}$ leads to (for $i, j, l > 1$):

$$d_{j1}(u) d_{11}(v) \equiv 0 \pmod{\mathcal{I}_2} \quad (3.24)$$

$$[d_{ij}(u), d_{11}(v)] \equiv 0 \pmod{\mathcal{I}_2} \quad (3.25)$$

$$[d_{ij}(u), d_{11}(v)] \equiv -\delta_{ij} \frac{\mathfrak{w}_{i1}(u, v) \bar{\mathfrak{w}}_{i1}(u, v)}{\mathfrak{b}(u, v) \bar{\mathfrak{b}}(u, v)} [d_{11}(u), d_{11}(v)] \pmod{\mathcal{I}_2}. \quad (3.26)$$

$$[d_{11}(u), d_{11}(v)] \equiv 0 \pmod{\mathcal{I}_2}. \quad (3.27)$$

Gathering all these equations, we get relation (3.20) for $k = 2$.

We now prove relation (3.23) for $k = 1, 2$. For $k = 1$, $d_{ij}^{(1)}(u) = d_{ij}(u)$, the ideal $\mathcal{I}_1$ is empty, and we have the starting algebra $\mathfrak{D}_{\mathbf{m}|\mathbf{n}}$, so that relation (3.23) for $k = 1$ just corresponds to the standard reflection equation. For $k = 2$, we use the following commutation relations:

$$[d_{ij}(u), d_{11}(v)] \equiv 0 \quad \text{for } i, j > 1 \quad \text{and} \quad [d_{11}(u), d_{11}(v)] \equiv 0, \quad (3.28)$$

$$d_{j1}(u) \equiv 0 \quad \text{for } j > 1 \pmod{\mathcal{I}_2}. \quad (3.29)$$

This implies that for $i, j, g, l > 1$, we have:

$$[d_{ij}^{(2)}(u), d_{gl}^{(2)}(v)] \equiv [d_{ij}(u^{(1)}), d_{gl}(v^{(1)})] \pmod{\mathcal{I}_2}. \quad (3.30)$$

Hence, it just remains to prove that the relation can be re-expressed in terms of $d_{rs}^{(1)}(u)$, $r, s > 1$, only. For such a purpose, we compute the commutation relations between $d_{i1}(u)$ and $d_{1j}(v)$:

$$\begin{aligned} d_{i1}(u) d_{1j}(v) &\equiv (-1)^{[j]([j]+[i])} \frac{\mathfrak{w}_{1j}(u, v)}{\mathfrak{b}(u, v)} d_{11}(v) d_{ij}(u) + \delta_{ij} (-1)^{[j]+[1]} \frac{\bar{\mathfrak{w}}_{i1}(u, v)}{\bar{\mathfrak{b}}(u, v)} d_{11}(v) d_{11}(u) \\ &\quad - (-1)^{[1]+[j][i]} \frac{\mathfrak{w}_{i1}(u, v)}{\mathfrak{b}(u, v)} d_{11}(u) d_{ij}(v) - \sum_{a=1}^{\mathbf{m}+\mathbf{n}} (-1)^{([i]+[j])[a]} \frac{\bar{\mathfrak{w}}_{1a}(u, v)}{\bar{\mathfrak{b}}(u, v)} d_{ia}(u) d_{aj}(v). \end{aligned} \quad (3.31)$$

Using this equation, one can compute⁴ for any polynomial function $f([a], [b], [c], [d])$:

$$\begin{aligned}
& \sum_{a=1}^{\mathbf{m}+\mathbf{n}} (-1)^{f([i],[l],[a],[g])} \frac{\bar{\mathbf{m}}_{ga}(u^{(1)}, v^{(1)})}{\bar{\mathbf{b}}(u^{(1)}, v^{(1)})} d_{ia}(u^{(1)}) d_{al}(v^{(1)}) \equiv \\
& \equiv \sum_{a=2}^{\mathbf{m}+\mathbf{n}} (-1)^{f([i],[l],[a],[g])} \frac{\bar{\mathbf{m}}_{ga}(u, v)}{\bar{\mathbf{b}}(u, v)} d_{ia}(u^{(1)}) d_{al}(v^{(1)}) \\
& - (-1)^{[1]+[g][i]+f([i],[l],[1],[g])} \frac{\bar{\mathbf{m}}_{i1}(u, v) \bar{\mathbf{m}}_{g1}(u^{(1)}, v^{(1)})}{\bar{\mathbf{b}}(u, v) \bar{\mathbf{a}}_1(u^{(1)}, v^{(1)})} d_{11}(u^{(1)}) d_{il}(v^{(1)}) \\
& + (-1)^{[l]([l]+[i])+f([i],[l],[1],[g])} \frac{\bar{\mathbf{m}}_{1l}(u, v) \bar{\mathbf{m}}_{g1}(u^{(1)}, v^{(1)})}{\bar{\mathbf{b}}(u, v) \bar{\mathbf{a}}_1(u^{(1)}, v^{(1)})} d_{11}(v^{(1)}) d_{il}(u^{(1)}) \\
& + \delta_{il} (-1)^{[1]+[i]+f([i],[l],[1],[g])} \frac{\bar{\mathbf{m}}_{i1}(u^{(1)}, v^{(1)}) \bar{\mathbf{m}}_{g1}(u^{(1)}, v^{(1)})}{\bar{\mathbf{b}}(u^{(1)}, v^{(1)}) \bar{\mathbf{a}}_1(u^{(1)}, v^{(1)})} d_{11}(u^{(1)}) d_{11}(v^{(1)}). \quad (3.32)
\end{aligned}$$

Now using (3.28), we can write the commutation relations for the new operator $d_{ij}^{(1)}(u)$ with (3.18). We extract the term $d_{i1}(u) d_{1j}(v)$ of each sum and using (3.32) we are left with only operators $d_{11}(u)$ and $d_{ij}(u)$ with $i, j \neq 1$. We use the transformation

$$d_{ij}^{(2)}(u) = d_{ij}^{(1)}(u^{(1)}) - \delta_{ij} \psi_1(u^{(1)}) d_{11}^{(1)}(u^{(1)}), \quad (3.33)$$

and we obtain the desired commutation relation plus some unwanted terms. It is a straightforward calculation to find that the unwanted terms cancel for all value of $i, j, g, l \in \{2, \dots, \mathbf{m} + \mathbf{n}\}$. This proves that $\hat{d}_{ij}(u)$ has the same commutation relation as (3.18) with a sum starting at 2 for $i, j > 1$. Thus, theorem 3.1 is proved for $k = 2$.

For $k > 2$, we first remark that one has $\mathcal{I}_k \subset \mathcal{I}_{k+1}$, so that one can use the results of step k in the proof of step $k + 1$. Then, the calculation becomes equivalent to the $k = 2$ case. The transformation of D operator for each step is:

$$D^{(k)}(u) = \mathbb{I}^{(k)} \hat{D}^{(k-1)}(u) \mathbb{I}^{(k)}, \quad k > 1 \quad (3.34)$$

$$\hat{D}^{(k)}(u) = \mathbb{I}^{(k)} \left(\hat{D}^{(k-1)}(u^{(k-1)}) - \psi_{k-1}(u^{(k-1)}) d_{kk}^{(k-1)}(u^{(k-1)}) \mathbb{I}^{(k)} \right) \mathbb{I}^{(k)}, \quad k > 1 \quad (3.35)$$

$$D^{(1)}(u) = \hat{D}^{(1)}(u) = \mathbb{I}^{(1)} D(u) \mathbb{I}^{(1)} = D(u) \quad (3.36)$$

A direct calculation using identity (B.10) gives the form (3.22). ■

Lemma 3.2 For $k = 1, 2, \dots, \mathbf{m} + \mathbf{n} - 1$, let $G^{(k)}$ be a linear combination of elements,

$$t_{i_1 j_1}(u_1) t'_{g_1 l_1}(\mathbf{u}(u_1)) \dots t_{i_p j_p}(u_p) t'_{g_p l_p}(\mathbf{u}(u_p)),$$

with all indices $i_r, j_r, g_r, l_r > k - 1$, and let $\mathcal{J}_k$ be the left ideal generated by $\{t_{ij}(u), t'_{ij}(u)\}$ for $i > j$ and $j < k$. Then, we have the following properties:

$$t_{ij}(u) G^{(k)} \equiv 0 \mod \mathcal{J}_k \quad \text{for } i > j \text{ and } j < k, \quad (3.37)$$

$$t'_{ij}(u) G^{(k)} \equiv 0 \mod \mathcal{J}_k \quad \text{for } i > j \text{ and } j < k, \quad (3.38)$$

$$[t_{ii}(u), G^{(k)}] \equiv 0 \mod \mathcal{J}_k \quad \text{for } i < k, \quad (3.39)$$

$$[t'_{ii}(u), G^{(k)}] \equiv 0 \mod \mathcal{J}_k \quad \text{for } i < k. \quad (3.40)$$

⁴To obtain this equation, we started from the lhs of (3.32), and changed the term $a = 1$ according to (3.31).

Moreover, the generators,

$$T^{(k)}(u) = \sum_{i,j=k}^{m+n} E_{ij} \otimes t_{ij}^{(k)}(u) \quad \text{and} \quad (T^{-1})^{(k)}(\mathbf{u}(u)) = \sum_{i,j=k}^{m+n} E_{ij} \otimes t'_{ij}{}^{(k)}(\mathbf{u}(u)), \quad (3.41)$$

$$t_{ij}^{(k)}(u) = t_{ij}(u^{(1\dots k-1)}) \quad \text{and} \quad t'_{ij}{}^{(k)}(\mathbf{u}(u)) = t'_{ij}(\mathbf{u}(u^{(1\dots k-1)})), \quad (3.42)$$

satisfy in $\mathcal{A}_{m|n}/\mathcal{J}_k$ the relation:

$$(T_2^{-1})^{(k)}(\mathbf{u}(u)) R_{12}^{(k)}(u, \mathbf{u}(u)) T_1^{(k)}(u) \equiv T_1^{(k)}(u) R_{12}^{(k)}(u, \mathbf{u}(u)) (T_2^{-1})^{(k)}(\mathbf{u}(u)) \mod \mathcal{J}_k. \quad (3.43)$$

Proof: As for theorem 3.1, the case $k = 1$ is just the definition of the algebra and relations (3.37)-(3.40) do not existe.

We prove the case $k = 2$, the proof for the other cases being similar. From the relation

$$T_2^{-1}(v) R_{12}(u, v) T_1(u) = T_1(u) R_{12}(u, v) T_2^{-1}(v) \quad (3.44)$$

one gets by projecting on $E_{ij} \otimes E_{gl}$:

$$\begin{aligned} \left[t'_{ij}(u), t_{gl}(v) \right] &= \delta_{il} \sum_{a=1}^{m+n} (-1)^{([j]+[a])([g]+[a])} \frac{\mathfrak{w}_{al}(v, u)}{\mathfrak{b}(v, u)} t_{ga}(v) t'_{aj}(u) \\ &\quad - \delta_{jg} \sum_{a=1}^{m+n} (-1)^{([g]+[a])([i]+[l])} \frac{\mathfrak{w}_{ga}(v, u)}{\mathfrak{b}(v, u)} t'_{ia}(u) t_{al}(v). \end{aligned} \quad (3.45)$$

From (3.45) we find for $i, j, g, l \neq 1$:

$$t'_{i1}(u) t_{11}(v) \equiv 0 \mod \mathcal{J}_2 \quad ; \quad \left[t'_{ij}(u), t_{g1}(v) \right] \equiv 0 \mod \mathcal{J}_2 \quad (3.46)$$

$$t_{g1}(v) t'_{11}(u) \equiv 0 \mod \mathcal{J}_2 \quad ; \quad \left[t'_{i1}(u), t_{gl}(v) \right] \equiv 0 \mod \mathcal{J}_2 \quad (3.47)$$

We also need the following commutation relation proved in [16]:

$$\left[t_{ij}(u), t_{k1}(v) \right] \equiv 0 \mod \mathcal{J}_2. \quad (3.48)$$

In the same way, one can compute:

$$\left[t'_{ij}(u), t'_{k1}(v) \right] \equiv 0 \mod \mathcal{J}_2. \quad (3.49)$$

Starting from the left-hand-side of relations (3.37)-(3.40), a recursive use of commutation relations (3.46)-(3.49) prove that one gets only terms with $t_{a1}(u)$ and $t'_{a1}(u)$ on the right, so that properties (3.37)-(3.40) hold for $k = 2$.

To prove (3.43), we start again with relation (3.45) with $i, j, g, l \neq 1$ and extract the first term in the summation:

$$\begin{aligned} \left[t'_{ij}(u), t_{gl}(v) \right] &= \delta_{il} \sum_{a=2}^{m+n} (-1)^{([j]+[a])([g]+[a])} \frac{\mathfrak{w}_{al}(v, u)}{\mathfrak{b}(v, u)} t_{ga}(v) t'_{aj}(u) \\ &\quad - \delta_{jg} \sum_{a=2}^{m+n} (-1)^{([g]+[a])([i]+[l])} \frac{\mathfrak{w}_{ga}(v, u)}{\mathfrak{b}(v, u)} t'_{ia}(u) t_{al}(v) \\ &\quad + \delta_{il} (-1)^{[j][g]+[1]} \frac{\mathfrak{w}_{1l}(v, u)}{\mathfrak{b}(v, u)} t_{g1}(v) t'_{1j}(u) - \delta_{jg} (-1)^{[g]([i]+[l])} \frac{\mathfrak{w}_{g1}(v, u)}{\mathfrak{b}(v, u)} t'_{i1}(u) t_{1l}(v). \end{aligned} \quad (3.50)$$

Inserting in this equation the relations (valid modulo $\mathcal{J}_2$):

$$t'_{i1}(u) t_{1l}(v) \equiv - \sum_{a=2}^{m+n} (-1)^{[a]([i]+[l])} \frac{\mathfrak{w}_{1a}(v, u)}{\mathfrak{a}_1(v, u)} t'_{ia}(u) t_{al}(v) + \delta_{il} \frac{\mathfrak{w}_{1l}(v, u)}{\mathfrak{a}_1(v, u)} t'_{11}(u) t_{11}(v), \quad (3.51)$$

$$\begin{aligned} (-1)^{[1]+[i][j]} t_{i1}(v) t'_{1j}(u) &\equiv - \sum_{a=2}^{m+n} (-1)^{([a]+[j])([a]+[i])} \frac{\mathfrak{w}_{a1}(v, u)}{\mathfrak{a}_1(v, u)} t_{ia}(v) t'_{aj}(u) \\ &\quad + \delta_{ji} \frac{\mathfrak{w}_{1l}(v, u)}{\mathfrak{a}_1(v, u)} t'_{11}(u) t_{11}(v), \end{aligned} \quad (3.52)$$

$$\left[t'_{11}(u), t_{11}(v) \right] \equiv 0, \quad (3.53)$$

and making the transformation $u \rightarrow \mathfrak{u}(u^{(1)})$ and $v \rightarrow u^{(1)}$ it is straightforward to end the proof. For $k > 2$ we use the same argument as in the proof of theorem 3.1. ■

Theorem 3.3 *In the coset $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}/\mathcal{J}_k \otimes \mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}/\mathcal{I}_k$, the coproduct takes the form*

$$\Delta(D_{[1]}^{(k)}(u)) \equiv T_{[2]}^{(k)}(u) D_{[1]}^{(k)}(u) (T_{[2]}^{-1})^{(k)}(\mathfrak{u}(u)) \mod \mathcal{J}_k, \quad (3.54)$$

where $[1]$ labels the space $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}/\mathcal{I}_k$, $[2]$ labels the space $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}/\mathcal{J}_k$ and Δ is the coproduct of $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$.

Proof: As in theorem 3.1, we just do the proof for $k = 2$, the other cases follow. From

$$\begin{aligned} \Delta\left(d_{ij}(u^{(1)}) - \delta_{ij} \psi_1(u^{(1)}) d_{11}(u^{(1)})\right) &= \sum_{a,b=1}^{m+n} \left\{ t_{ia}(u^{(1)}) t'_{bj}(\mathfrak{u}(u^{(1)})) \otimes d_{ab}(u^{(1)}) \right. \\ &\quad \left. - \delta_{ij} \psi_1(u^{(1)}) t_{1a}(u^{(1)}) t'_{b1}(\mathfrak{u}(u^{(1)})) \otimes d_{ab}(u^{(1)}) \right\}. \end{aligned}$$

and using the quotient $\mathcal{I}_2$ and $\mathcal{J}_2$, it follows:

$$\begin{aligned} \Delta\left(d_{ij}(u^{(1)}) - \delta_{ij} \psi_1(u^{(1)}) d_{11}(u^{(1)})\right) &= \sum_{a,b=2}^{m+n} t_{ia}(u^{(1)}) t'_{bj}(\mathfrak{u}(u^{(1)})) \otimes d_{ab}(u^{(1)}) + \\ &\quad + t_{i1}(u^{(1)}) t'_{1j}(\mathfrak{u}(u^{(1)})) \otimes d_{11}(u^{(1)}) - \delta_{ij} \psi_1(u^{(1)}) t_{11}(u^{(1)}) t'_{11}(\mathfrak{u}(u^{(1)})) \otimes d_{11}(u^{(1)}). \end{aligned}$$

Using the commutation relation (3.45) for the second term we find:

$$\Delta\left(d_{ij}(u^{(1)}) - \delta_{ij} \psi_1(u^{(1)}) d_{11}(u^{(1)})\right) = \sum_{a,b=2}^{m+n} t_{ia}(u^{(1)}) t'_{bj}(\mathfrak{u}(u^{(1)})) \otimes \left(d_{ij}(u^{(1)}) - \delta_{ij} \psi_1(u^{(1)}) d_{11}(u^{(1)})\right).$$

Theorem 3.1 and lemma 3.2 allow us to generalise this result to each k . ■

4 Highest weight representations

The fundamental point in using the ABA is to know a pseudo-vacuum for the model. In the mathematical framework it is equivalent to know a highest weight representation for the algebra which underlies the model. Since the generators of the algebra $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ can be constructed from the $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ ones, see eq. (3.10), we first describe how to construct highest representations for the infinite dimensional (graded) algebras $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ from highest weight representation of the finite dimensional Lie subalgebras $gl(\mathfrak{m}|\mathfrak{n})$ or $\mathcal{U}_q(\mathfrak{m}|\mathfrak{n})$. Next, we show how these representations induce (for diagonal $K^-(u)$ matrix) a representation for $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ with same highest weight vector.

4.1 Finite dimensional representations of $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$

Definition 4.1 A representation of $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ is called *highest weight* if there exists a nonzero vector Ω such that,

$$t_{ii}(u)\Omega = \lambda_i(u)\Omega \quad \text{and} \quad t_{ij}(u)\Omega = 0 \quad \text{for } i > j, \quad (4.1)$$

for some scalars $\lambda_i(u) \in \mathbb{C}[[u^{-1}]]$. $\lambda(u) = (\lambda_1(u), \dots, \lambda_{\mathfrak{m}+\mathfrak{n}}(u))$ is called the *highest weight* and Ω the *highest weight vector*.

It is known (see [27–30]) that any finite-dimensional irreducible representation of $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ is highest weight and that it contains a unique (up to scalar multiples) highest weight vector. To construct such representations, one uses the evaluation morphism, which relates the infinite dimensional algebra $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ to its finite dimensional subalgebra $\mathcal{B}_{\mathfrak{m}|\mathfrak{n}}$ (see [16]). From the evaluation morphism ev_a (with $a \in \mathbb{C}$) and a highest weight representation π_μ of $\mathcal{B}_{\mathfrak{m}|\mathfrak{n}}$ (where μ is a $\mathcal{B}_{\mathfrak{m}|\mathfrak{n}}$ highest weight), one can construct a highest weight representation of $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$, called evaluation representation:

$$\rho_a^\mu = ev_a \circ \pi_\mu : \mathcal{A}_{\mathfrak{m}|\mathfrak{n}} \xrightarrow{ev_a} \mathcal{B}_{\mathfrak{m}|\mathfrak{n}} \xrightarrow{\pi_\mu} \mathcal{V}_\lambda. \quad (4.2)$$

The weight of this evaluation representation is given by $\lambda(u) = (\lambda_1(u), \dots, \lambda_{\mathfrak{m}+\mathfrak{n}}(u))$, with

$$\lambda_j(u) = \begin{cases} u - a - (-1)^{[j]} \hbar \mu_j & \text{for } \mathcal{Y}(\mathfrak{m}|\mathfrak{n}) \\ (-1)^{[j]} \left(\frac{u}{a} \eta_j q^{\mu_j} - \frac{a}{u} \eta_j q^{-\mu_j} \right) & \text{for } \widehat{\mathcal{U}}_q(\mathfrak{m}|\mathfrak{n}) \end{cases} \quad j = 1, \dots, \mathfrak{m} + \mathfrak{n}, \quad (4.3)$$

where μ_j , $j = 1, \dots, \mathfrak{m} + \mathfrak{n}$ are the weights of the $\mathcal{B}_{\mathfrak{m}|\mathfrak{n}}$ representation. More generally, one constructs tensor product of evaluation representations using the coproduct of $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$,

$$\left(\bigotimes_{i=1}^L \rho_{a_i}^{\mu^{(i)}} \right) \circ \Delta^{(L)}(T(u)) = \rho_{a_1}^{\mu^{(1)}}(T(u)) \dot{\otimes} \rho_{a_2}^{\mu^{(2)}}(T(u)) \dot{\otimes} \dots \dot{\otimes} \rho_{a_L}^{\mu^{(L)}}(T(u)), \quad (4.4)$$

where $\mu^{(i)} = (\mu_1^{(i)}, \dots, \mu_{\mathfrak{m}+\mathfrak{n}}^{(i)})$, $i = 1, \dots, L$, are the weights of the $\mathcal{B}_{\mathfrak{m}|\mathfrak{n}}$ representations. This provides a $\mathcal{A}_{\mathfrak{m}|\mathfrak{n}}$ representation with weight,

$$\lambda_j(u) = \prod_{i=1}^L \lambda_j^{(i)}(u), \quad j = 1, \dots, \mathfrak{m} + \mathfrak{n}, \quad (4.5)$$

where $\lambda_j^{(i)}(u)$ have the form (4.3).

4.2 Representations of $T^{-1}(u)$ from $T(u)$

The construction of the finite dimensional representations for $T^{-1}(u)$ in relation with the $T(u)$ ones is different for the $gl(\mathfrak{n})$ and the super symmetric case $gl(\mathfrak{m}|\mathfrak{n})$. For the $gl(\mathfrak{n})$ algebra, the representations are constructed using the quantum determinant $qdet(T(u))$ and the comatrix $\widehat{T}(u)$ see [31], while for the $gl(\mathfrak{m}|\mathfrak{n})$ superalgebra, one uses the Liouville contraction, the quantum Berezinian $Ber(T(u))$ [32] and crossing symmetry of $T(u)$.

We define for this section:

$$u_{\{k\}} = \begin{cases} u + \hbar k \\ u q^{-k} \end{cases} \quad \text{and} \quad f_{ij}(\sigma) = \begin{cases} (-1)^{i+j+1} s(\sigma) & \text{for } \mathcal{Y}(\mathfrak{n}) \text{ and } \mathcal{Y}(\mathfrak{m}|\mathfrak{n}) \\ (-q)^{l(\sigma)+i-j} & \text{for } \widehat{\mathcal{U}}_q(\mathfrak{n}) \text{ and } \widehat{\mathcal{U}}_q(\mathfrak{m}|\mathfrak{n}) \end{cases} \quad (4.6)$$

where $s(\sigma)$ is the sign of the permutation σ and $l(\sigma)$ its length.

$\mathcal{A}_n$ case: We use the $\mathcal{A}_n$ quantum determinant $qdet(T(u))$ which generates the center of $\mathcal{A}_n$,

$$qdet(T(u)) = \sum_{\sigma \in S_n} f_{00}(\sigma) \prod_{i=1}^n t_{i\sigma(i)}(u_{\{i-n\}}), \quad (4.7)$$

and the quantum comatrix,

$$\widehat{T}(u) = \sum_{ij=1} E_{ij} \otimes \widehat{t}_{ij}(u) \quad (4.8)$$

$$\begin{aligned} \widehat{t}_{ij}(u) &= \sum_{\sigma \in S_n} f_{ij}(\sigma)^{-1} t_{1a_{\sigma(1)}}(u_{\{2-n\}}) \cdots t_{i-1a_{\sigma(i-1)}}(u_{\{i-n\}}) t_{i+1a_{\sigma(i)}}(u_{\{i+1-n\}}) \cdots t_{na_{\sigma(n)}}(u) \\ &\text{with } (a_1, \dots, a_{n-1}) = (1, \dots, j-1, j+1, \dots, n) \end{aligned} \quad (4.9)$$

which obeys $\widehat{T}(u) T(u_{\{1-n\}}) = qdet(T(u))$. This equation allows to relate $T^{-1}(u)$ to $\widehat{T}(u)$:

$$T^{-1}(u) = \sum_{i,j=1}^{m+n} E_{ij} \otimes t'_{ij}(u) = \frac{\widehat{T}(u_{\{n-1\}})}{qdet(T(u_{\{n-1\}}))}. \quad (4.10)$$

To write the form of the highest weight irreducible representation for $T^{-1}(u)$, one first computes the action of $qdet(T(u))$ and $\widehat{t}_{ii}(u)$ on Ω :

$$qdet(T(u)) \Omega = \prod_{i=1}^n \lambda_i(u_{\{i-n\}}) \Omega, \quad (4.11)$$

$$\widehat{t}_{ii}(u) \Omega = \lambda_1(u_{\{2-n\}}) \cdots \lambda_{i-1}(u_{\{i-n\}}) \lambda_{i+1}(u_{\{i+1-n\}}) \cdots \lambda_n(u) \Omega. \quad (4.12)$$

Then, since $\widehat{t}_{ij}(u) \Omega = 0$ for $i > j$, one finds:

$$\begin{aligned} t'_{ii}(u) \Omega &= \lambda'_i(u) \Omega \quad \text{with} \quad \lambda'_i(u) = \left(\prod_{k=1}^{i-1} \frac{\lambda_k(u_{\{k\}})}{\lambda_k(u_{\{k-1\}})} \right) \frac{1}{\lambda_i(u_{\{i-1\}})}, \\ t'_{ij}(u) \Omega &= 0 \quad \text{if } i > j. \end{aligned} \quad (4.13)$$

$\mathcal{A}_{m|n}$ case: First, one has to prove that Ω is a highest weight vector of $T^{-1}(u)$. The proof is done in [15] for the super-Yangian case. The quantum superalgebra $\widehat{\mathcal{U}}_q(\mathfrak{m}|n)$ case is done in the following theorem:

Theorem 4.2 *For the quantum superalgebra $\widehat{\mathcal{U}}_q(\mathfrak{m}|n)$, the highest weight vector Ω of $T(u)$ is also a highest weight vector of $T^{-1}(u)$.*

$$t'_{ii}(u) \Omega = \lambda'_i(u) \Omega \quad \text{and} \quad t'_{ij}(u) \Omega = 0 \quad \text{if } i > j \quad (4.14)$$

Proof: To prove this theorem we must use the commutation relation between the modes of $T(u) = L^+(u)$ and $T^{-1}(u) = (L^+)^{-1}(u) = \sum_{i,j=1}^{m+n} \bar{L}_{ij}^+(u)$. As $L^+(u)$ is a formal Taylor series in u , its inverse is also a formal Taylor series of u :

$$L_{ij}^+(u) = \sum_{n=0}^{\infty} L_{ij}^{(n)} u^{2n} \quad \text{and} \quad \bar{L}_{ij}^+(u) = \sum_{n=0}^{\infty} \bar{L}_{ij}^{(n)} u^{2n}. \quad (4.15)$$

Projecting the commutation relation (3.45) on the modes we find the following relation:

$$\begin{aligned}
[\bar{L}_{ij}^{(p)}; L_{kl}^{(q)}] &= \delta_{il} \sum_{a=1}^{m+n} (-1)^{([a]+[j])([a]+[k])} \left(c_{al}^+ \sum_{b=0}^p L_{ka}^{(q+b)} \bar{L}_{aj}^{(p-b)} - c_{al}^- \sum_{b=1}^p L_{ka}^{(q+b)} \bar{L}_{aj}^{(p-b)} \right) \\
&- \delta_{jk} \sum_{a=1}^{m+n} (-1)^{([i]+[l])([a]+[k])} \left(c_{ka}^+ \sum_{b=0}^p \bar{L}_{ia}^{(p-b)} L_{al}^{(q+b)} - c_{ka}^- \sum_{b=1}^p \bar{L}_{ia}^{(p-b)} L_{al}^{(q+b)} \right), \\
\text{with } c_{al}^\pm &= q^{\pm(1-2[l])} - q^{\text{sign}(l-a)(1-2[l])}.
\end{aligned} \tag{4.16}$$

From the relation $T(u) T^{-1}(u) = T^{-1}(u) T(u) = \mathbb{I}$ we find:

$$\sum_{a=1}^{m+n} \sum_{q=0}^p (-1)^{([i]+[a])([a]+[j])} \bar{L}_{ia}^{(q)} L_{aj}^{(p-q)} = \sum_{a=1}^{m+n} \sum_{q=0}^p (-1)^{([i]+[a])([a]+[j])} L_{ia}^{(q)} \bar{L}_{aj}^{(p-q)} = \delta_{ij} \delta_{0p}. \tag{4.17}$$

We also know that $L_{ij}^{(0)} = 0$ for $i < j$ and that $L_{ii}^{(0)}$ has an inverse. First we prove the same properties for $\bar{L}^{(0)}$. From (4.17), we deduce:

$$\sum_{a=1}^{m+n} (-1)^{([i]+[a])([a]+[j])} L_{ia}^{(0)} \bar{L}_{aj}^{(0)} = 0 \quad \text{for } i \neq j. \tag{4.18}$$

As $L_{ii}^{(0)} \neq 0$, we find for $i = 1$, $\bar{L}_{1j}^{(0)} = 0$. By induction on i , we find $\bar{L}_{ij}^{(0)} = 0$ for $i < j$. Then,

$$\sum_{a=1}^{m+n} (-1)^{[i]+[a]} L_{ia}^{(0)} \bar{L}_{ai}^{(0)} = \sum_{a=1}^{m+n} (-1)^{[i]+[a]} \bar{L}_{ia}^{(0)} L_{ai}^{(0)} = 1 \tag{4.19}$$

implies that $L_{ii}^{(0)} \bar{L}_{ii}^{(0)} = \bar{L}_{ii}^{(0)} L_{ii}^{(0)} = 1$.

Now, we have to prove that Ω is a highest weight vector of $\bar{L}_{ij}^{(p)}$. We already know that:

$$L_{ij}^{(p)} \Omega = 0 \quad \text{for } i < j \quad \text{and} \quad L_{ii}^{(p)} \Omega = \lambda_i^{(p)} \Omega. \tag{4.20}$$

We can write from (4.17) with $i > j$:

$$\sum_{q=0}^p \lambda_j^{(p-q)} \bar{L}_{ij}^{(q)} \Omega = - \sum_{a=1}^{j-1} \sum_{q=0}^{p-1} (-1)^{([k]+[a])([i]+[j])} \bar{L}_{ia}^{(q)} L_{aj}^{(p-q)} \Omega. \tag{4.21}$$

To prove that $L_{ij}^{(p)} \Omega = 0$ for $i < j$, we use a double induction, on p and on i . We already proved directly that $\bar{L}_{ij}^{(0)} \Omega = 0$, $i < j$. For $p = 1$ we have:

$$\lambda_j^{(0)} \bar{L}_{ij}^{(1)} \Omega = - \sum_{a=1}^{j-1} (-1)^{([k]+[a])([i]+[j])} \bar{L}_{ia}^{(0)} L_{aj}^{(1)} \Omega = - \sum_{a=1}^{j-1} (-1)^{([k]+[a])([i]+[j])} [\bar{L}_{ia}^{(0)}, L_{aj}^{(1)}] \Omega.$$

Using the commutation relations (4.16) we find:

$$[\bar{L}_{ia}^{(0)}, L_{aj}^{(1)}] \Omega = -q^{-1+2[a]} (q - q^{-1}) (-1)^{([i]+[j])[a]} \sum_{b=1}^{a-1} (-1)^{[b]([i]+[b]+[j])} [\bar{L}_{ib}^{(0)}, L_{bj}^{(1)}] \Omega. \tag{4.22}$$

We get a triangular system in a , so that the property is proved for $p = 1$ by induction on a . For a general p we use the same method.

Finally, we prove that $\bar{L}_{ii}^{(p)} \Omega = \bar{\lambda}_i^{(p)} \Omega$. For $p = 0$, from the equation $\bar{L}_{ii}^{(0)} L_{ii}^{(0)} = 1$, we already know that $\bar{L}_{ii}^{(0)} \Omega = (\lambda_i^{(0)})^{-1} \Omega$. We prove the property of the general case by a double induction, assuming the property is true for $p - 1$ and starting from:

$$\sum_{q=0}^p \lambda_i^{(p-q)} \bar{L}_{ii}^{(q)} \Omega = \left(\delta_{0,p} - \sum_{a=1}^{i-1} \sum_{q=0}^{p-1} (-1)^{[i]+[a]} [\bar{L}_{ia}^{(q)}, L_{ai}^{(p-q)}] \right) \Omega. \quad (4.23)$$

From the commutation relation we get:

$$\begin{aligned} [\bar{L}_{ia}^{(q)}, L_{ai}^{(p-q)}] \Omega &= \sum_{b=1}^a \left(c_{bi}^- \sum_{r=1}^q [\bar{L}_{ib}^{(q-r)}, L_{bi}^{(p-q+r)}] - c_{ab}^+ \sum_{r=0}^q [\bar{L}_{ib}^{(q-r)}, L_{bi}^{(p-q+r)}] \right) \Omega, \\ [\bar{L}_{ia}^{(0)}, L_{ai}^{(q)}] \Omega &= -q^{-1+2[a]} (q - q^{-1}) (-1)^{[a]} \sum_{b=1}^{a-1} [\bar{L}_{ib}^{(0)}, L_{bi}^{(q)}] \Omega = 0. \end{aligned} \quad (4.24)$$

From the second equation, equals to zero by induction on a , we find:

$$[\bar{L}_{ia}^{(q)}, L_{ai}^{(p-q)}] \Omega = -q^{-1+2[a]} (q - q^{-1}) (-1)^{[a]} \sum_{b=1}^{a-1} \sum_{r=1}^q [\bar{L}_{ib}^{(q-r)}, L_{bi}^{(p-q+r)}] \Omega. \quad (4.25)$$

By induction on q then on a we find the last equality equals to zero. Thus, we have the relation:

$$\sum_{q=0}^p \lambda_i^{(p-q)} \bar{L}_{ii}^{(q)} \Omega = \delta_{0,p} \Omega \quad (4.26)$$

It follows that $\bar{L}_{ii}^{(q)} \Omega = \bar{\lambda}_i^{(q)} \Omega$ with $\bar{\lambda}_i^{(q)}$ rational function of $\lambda_i^{(q)}, \dots, \lambda_i^{(0)}$. ■

Secondly, we use the crossing symmetry of the monodromy matrix $T_a(u)$ and the quantum Berezinian to give an explicit expression of the weight of $T^{-1}(u)$. Let us introduce:

$$T^*(u) = (T^{-1}(u))^t. \quad (4.27)$$

The crossing symmetry takes the form for $\mathcal{A}_{m|n}$ (see [15, 33]):

$$(T^t(u))^{-1} = \frac{1}{Z(u_{\{n-m\}})} M U T^*(u_{\{n-m\}}) U M^{-1} \quad \text{with} \quad U = \sum_{i=1}^{m+n} (-1)^{[i]} E_{ii}, \quad (4.28)$$

where the Liouville contraction $Z(u)$ lies in the centre of $\mathcal{A}_{m|n}$. It can be written in term of the Berezinian, that itself relies on the quantum determinant (4.7):

$$Ber(T(u)) = qdet\left(T^{(m)}(u_{\{m-n-1\}})\right) \widetilde{qdet}\left((T^*)^{(n)}(u_{\{n\}})\right) = \prod_{i=1}^m \lambda_i(u_{\{i-n-1\}}) \prod_{i=1}^n \lambda'_{i+m}(u_{\{i-n-1\}}) \quad (4.29)$$

$$Z(u) = \frac{Ber(T(u_{\{1\}}))}{Ber(T(u))} \quad \text{with} \quad \widetilde{qdet}(T^{(n)}(u)) = \sum_{\sigma \in S_n} f_{00}(\sigma)^{-1} \prod_{i=1}^n t_{i+m, \sigma(i)+m}(u_{\{n-i\}}), \quad (4.30)$$

with $T^{(k)}(u) = \mathbb{I}^{(k)} T(u) \mathbb{I}^{(k)}$, $\forall k$ and $\mathbb{I}^{(k)}$ defined in (2.14). In $\mathcal{A}_{0|n} \subset \mathcal{A}_{m|n}$, we have:

$$(T^{(n)}(u)^t)^{-1} = z(u) M T^{(n)*}(u_{\{n\}}) M^{-1}, \quad (4.31)$$

where $z(u)$, the $\mathcal{A}_{0|n}$ Liouville contraction, can be written in term of the quantum determinant:

$$z(u) = \frac{\widetilde{qdet}(T^{(n)}(u_{\{1\}}))}{\widetilde{qdet}(T^{(n)}(u))} = \prod_{i=1}^n \frac{\lambda_{i+m}(u_{\{n-i+1\}})}{\lambda_{i+m}(u_{\{n-i\}})}. \quad (4.32)$$

Lemma 4.3 *For the superalgebra $\mathcal{A}_{m|n}$, we have:*

$$\begin{aligned} t'_{ii}(u) \Omega &= \lambda'_i(u) \Omega \quad \text{and} \quad t'_{ij}(u) \Omega = 0 \quad \text{for } i > j, \\ \text{with } \lambda'_i(u) &= \begin{cases} \frac{1}{\lambda_i(u_{\{i-1\}})} \left(\prod_{k=1}^{i-1} \frac{\lambda_k(u_{\{k\}})}{\lambda_k(u_{\{k-1\}})} \right) & \text{for } 1 \leq i \leq m \\ \frac{Z(u)}{\lambda_i(u_{\{2m-i\}})} \left(\prod_{k=i+1}^{m+n} \frac{\lambda_k(u_{\{2m+1-k\}})}{\lambda_k(u_{\{2m-k\}})} \right) & \text{for } m+1 \leq i \leq m+n \end{cases} \end{aligned} \quad (4.33)$$

Proof: From theorem 4.2, we have,

$$T^{-1}(u) \Omega = \begin{pmatrix} T^{-1}(u)^{(m)} & * \\ 0 & T^{-1}(u)^{(n)} \end{pmatrix} \Omega, \quad (4.34)$$

$$T^*(u) \Omega = \begin{pmatrix} T^*(u)^{(m)} & 0 \\ * & T^*(u)^{(n)} \end{pmatrix} \Omega. \quad (4.35)$$

Multiplying (4.34) by $T(u)$ and (4.35) by $UM^{-1}T^t(u_{\{n-m\}})MU$, one gets:

$$T^{(m)}(u) (T^{-1})^{(m)}(u) \Omega = \Omega \quad \text{and} \quad (M^{-1})^{(n)} (T^t)^{(n)}(u_{\{n-m\}}) M^{(n)} (T^*)^{(n)}(u) \Omega = Z(u) \Omega.$$

Finally, upon multiplication by $T^{(m)}(u)^{-1}$ and $(T^t)^{(n)}(u)^{-1}$, one is led to

$$(T^{-1})^{(m)}(u) \Omega = (T^{(m)})^{-1}(u) \Omega \quad \text{and} \quad (T^*)^{(n)}(u) \Omega = Z(u) z(u_{\{m-n\}}) (T^{(n)})^*(u_{\{m\}}) \Omega \quad (4.36)$$

that gives the lemma. ■

4.3 Finite dimensional representations of $\mathfrak{D}_{m|n}$ from $\mathcal{A}_{m|n}$ ones

For the study of the representations of the reflection algebra, we follow essentially the lines given in [26] for the reflection algebra based on the Yangian of $gl(n)$ and in [15] for the reflection algebra based on the super-Yangian of $gl(m|n)$.

Theorem 4.4 *If Ω is a highest weight vector of $\mathcal{A}_{m|n}$, with eigenvalue $(\lambda_1(u), \dots, \lambda_{m+n}(u))$, then, when $K^-(z) = \text{diag}(\kappa_1(u), \dots, \kappa_{m+n}(u))$, Ω is also a highest weight vector for $\mathfrak{D}_{m|n}$,*

$$d_{ij}(u) \Omega = 0 \quad \text{for } i > j, \quad \text{and} \quad d_{ii}(u) \Omega = \Lambda_i(u) \Omega, \quad (4.37)$$

with eigenvalues:

$$\Lambda_i(u) = \mathcal{K}_i(u) \lambda_i(u) \lambda'_i(\mathbf{u}(u)) + \sum_{k=1}^{i-1} \psi_k(u^{\overline{(1\dots k-1)}}) \mathcal{K}_k(u) \lambda_k(u) \lambda'_k(\mathbf{u}(u)), \quad (4.38)$$

$$\mathcal{K}_i(u) = \kappa_i(u) - \sum_{k=1}^{i-1} \kappa_k(u) \frac{\mathfrak{w}_{ik}(u^{\overline{(1\dots k-1)}}, \mathbf{u}(u^{\overline{(1\dots k-1)}}))}{\mathfrak{a}_{i-1}(u^{\overline{(1\dots i-2)}}, \mathbf{u}(u^{\overline{(1\dots i-2)}}))} q^{i-k-1-2\sum_{l=k+1}^{i-1} [l]}. \quad (4.39)$$

Proof: First, we prove $d_{ij}(u) \Omega = 0$ for $j < i$. One computes:

$$\begin{aligned} d_{ij}(u) \Omega &= \sum_{a=1}^{j-1} (-1)^{([i]+[a])([a]+[j])} \kappa_a(u) t_{ia}(u) t'_{aj}(\mathbf{u}(u)) \Omega \\ &= - \sum_{a=1}^{j-1} \kappa_a(u) \left[t'_{aj}(\mathbf{u}(u)), t_{ia}(u) \right] \Omega. \end{aligned} \quad (4.40)$$

Applying the super-commutator on Ω with the constraint $a \leq j < i$, one gets:

$$\left[t'_{aj}(u), t_{ia}(\mathbf{u}(u)) \right] \Omega = - \sum_{b=1}^{j-1} \frac{\mathfrak{w}_{ba}(u, \mathbf{u}(u))}{\mathfrak{b}(u, \mathbf{u}(u))} \left[t'_{bj}(\mathbf{u}(u)), t_{ib}(u) \right] \Omega. \quad (4.41)$$

Considering the case $a = j$, one obtains:

$$\sum_{b=1}^{j-1} \left[t'_{bj}(\mathbf{u}(u)), t_{ib}(u) \right] \Omega = 0. \quad (4.42)$$

Plugging this result in the former equation, we get :

$$\left[t'_{aj}(u), t_{ia}(\mathbf{u}(u)) \right] \Omega = \frac{\mathfrak{w}_{a-1,a}(u, \mathbf{u}(u)) - \mathfrak{w}_{a+1,a}(u, \mathbf{u}(u))}{\mathfrak{a}_a(u, \mathbf{u}(u)) - \mathfrak{w}_{a-1,a}(u, \mathbf{u}(u))} \sum_{b=a+1}^{j-1} \left[t'_{bj}(\mathbf{u}(u)), t_{ib}(u) \right] \Omega. \quad (4.43)$$

By iteration ($a = j-1, \dots, a = 1$) one finds:

$$\left[t'_{aj}(u), t_{ia}(\mathbf{u}(u)) \right] \Omega = 0, \quad (4.44)$$

which proves that $d_{ij}(u) \omega = 0$, $j < i$.

Secondly, we prove $d_{ii}(u) \Omega = \Lambda_i(u) \Omega$. Acting on Ω with $d_{ii}(u)$ one obtains:

$$d_{ii}(u) \Omega = \kappa_i(u) \lambda_i(u) \lambda'_i(\mathbf{u}(u)) \Omega + \sum_{a=1}^{i-1} (-1)^{[i]+[a]} \kappa_a(u) t_{ia}(u) t'_{ai}(\mathbf{u}(u)) \Omega. \quad (4.45)$$

One can restrict this problem to the computation of $t_{ia}(u) t'_{ai}(\mathbf{u}(u)) \Omega$ for $i > a$ in term of the eigenvalues $\lambda_i(u) \lambda'_i(\mathbf{u}(u))$. From the relation (3.45), we get

$$\begin{aligned} (-1)^{[i]+[a]} t_{ia}(u) t'_{ai}(\mathbf{u}(u)) \Omega &= - \sum_{b=1}^i (-1)^{[i]+[b]} \frac{\mathfrak{w}_{ba}(u, \mathbf{u}(u))}{\mathfrak{b}(u, \mathbf{u}(u))} t_{ib}(u) t'_{bi}(\mathbf{u}(u)) \Omega \\ &\quad + \sum_{b=1}^a \frac{\mathfrak{w}_{ib}(u, \mathbf{u}(u))}{\mathfrak{b}(u, \mathbf{u}(u))} t'_{ab}(\mathbf{u}(u)) t_{ba}(u) \Omega. \end{aligned} \quad (4.46)$$

Applying (3.45) on Ω for $i = j = k = l$, one finds the identity:

$$\sum_{b=1}^{i-1} \frac{\mathfrak{w}_{ib}(u, \boldsymbol{\iota}(u))}{\mathfrak{b}(u, \boldsymbol{\iota}(u))} t'_{ib}(\boldsymbol{\iota}(u)) t_{bi}(u) \Omega = \sum_{b=1}^{i-1} (-1)^{[i]+[b]} \frac{\mathfrak{w}_{bi}(u, \boldsymbol{\iota}(u))}{\mathfrak{b}(u, \boldsymbol{\iota}(u))} t_{ib}(u) t'_{bi}(\boldsymbol{\iota}(u)) \Omega.$$

Let $F_{ij} = t_{ij}(u) t'_{ji}(\boldsymbol{\iota}(u)) \Omega$. Using the two previous equations, one finds for $j < i$:

$$\begin{aligned} F_{ij} &= (-1)^{[i]+[j]} \frac{\mathfrak{w}_{ij}(u, \boldsymbol{\iota}(u))}{\mathfrak{b}(u, \boldsymbol{\iota}(u))} (F_{jj} - F_{ii}) + \sum_{a=1}^{j-1} (-1)^{[i]+[a]} \frac{\mathfrak{w}_{ai}(u, \boldsymbol{\iota}(u))}{\mathfrak{b}(u, \boldsymbol{\iota}(u))} F_{ja} \\ &\quad - \sum_{a=1}^{i-1} (-1)^{[j]+[a]} \frac{\mathfrak{w}_{aj}(u, \boldsymbol{\iota}(u))}{\mathfrak{b}(u, \boldsymbol{\iota}(u))} F_{ia}. \end{aligned} \quad (4.47)$$

It is then easy (but lengthy) to show that the solution is:

$$\begin{aligned} F_{ij} &= (-1)^{[i]+[j]} \mathfrak{m}_{ij}(u^{\overline{(1\dots j-1)}}, \boldsymbol{\iota}(u^{\overline{(1\dots j-1)}})) \left[\frac{F_{jj}}{\mathfrak{a}_j(u^{\overline{(1\dots j-1)}}, \boldsymbol{\iota}(u^{\overline{(1\dots j-1)}}))} \right. \\ &\quad - \frac{q^{(i-j-1-2\sum_{a=j+1}^{i-1}[a])}}{\mathfrak{a}_{i-1}(u^{\overline{(1\dots i-2)}}, \boldsymbol{\iota}(u^{\overline{(1\dots i-2)}}))} F_{ii} \\ &\quad \left. - \sum_{a=j+1}^{i-1} \frac{\mathfrak{m}_{ia}(u^{\overline{(1\dots a-1)}}, \boldsymbol{\iota}(u^{\overline{(1\dots a-1)}})) q^{(a-j-1-2\sum_{l=j+1}^{a-1}[l])}}{\mathfrak{a}_j(u^{\overline{(1\dots a-1)}}, \boldsymbol{\iota}(u^{\overline{(1\dots a-1)}})) \mathfrak{a}_{a-1}(u^{\overline{(1\dots a-2)}}, \boldsymbol{\iota}(u^{\overline{(1\dots a-2)}}))} F_{aa} \right]. \end{aligned} \quad (4.48)$$

One must use relations (B.7)-(B.9) between functions. Plugging the value of F_{ik} into the equation (4.45), after some rearrangement one gets the eigenvalues $\Lambda_i(u)$. $\blacksquare$

5 Algebraic Bethe ansatz for $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ with $\mathfrak{m} + \mathfrak{n} = 2$

In this section, we remind the framework of the Algebraic Bethe Ansatz (ABA) [34] introduced in order to compute transfer matrix eigenvalues and eigenvectors. For $\mathfrak{m} + \mathfrak{n} = 2$, one can consider three different algebras: $\mathfrak{D}_{0|2}$, $\mathfrak{D}_{2|0}$ and $\mathfrak{D}_{1|1}$. The method follows the same steps as the closed chain case, up to a preliminary step. We write the monodromy matrix in the following matricial form:

$$D(u) = \begin{pmatrix} d_{11}(u) & d_{12}(u) \\ d_{21}(u) & d_{22}(u) \end{pmatrix}. \quad (5.1)$$

In the open case the transfer matrix have the form:

$$d(u) = \text{str}(K_a^+(u) D_a(u)) = (-1)^{[1]} m_1 k(u) d_{11}(u) + (-1)^{[2]} m_2 d_{22}(u), \quad (5.2)$$

$$K^+(u) = M K(u). \quad (5.3)$$

The matrix K is construct from the solution (3.9):

$$K(u) = \begin{cases} \mathbb{I} & \text{for } a_+ = 0 \\ \text{diag}(k(u), 1) & \text{for } a_+ = 1 \end{cases} \quad \text{with } k(u) = \begin{cases} \frac{-u - \frac{\mathfrak{m}-\mathfrak{n}}{2}\hbar - c_+}{u + \frac{\mathfrak{m}-\mathfrak{n}}{2}\hbar - c_+} & \text{for } \mathcal{Y}(\mathfrak{m}|\mathfrak{n}) \\ \frac{u^{-2} q^{-\frac{\mathfrak{m}-\mathfrak{n}}{2}} - c_+^2}{u^2 q^{\frac{\mathfrak{m}-\mathfrak{n}}{2}} - c_+^2} & \text{for } \widehat{\mathcal{U}}_q(\mathfrak{m}|\mathfrak{n}) \end{cases}. \quad (5.4)$$

Remark that for the particular case $\mathbf{m} + \mathbf{n} = 2$, the form chosen for $K^+(u)$ exhausts all possible diagonal solutions. We recall that for $K^-(u)$ we keep the general diagonal solution (3.9). Let Ω be the pseudo-vacuum state:

$$d_{11}(u) \Omega = \Lambda_1(u) \Omega, \quad d_{22}(u) \Omega = \Lambda_2(u) \Omega, \quad d_{21}(u) \Omega = 0. \quad (5.5)$$

Looking at the commutation relations (3.18) for $\mathbf{m} + \mathbf{n} = 2$, one can see that the $d_{22}(u) d_{12}(v)$ exchange relation is not symmetric to the $d_{11}(u) d_{12}(v)$ one. In order to compensate this asymmetry, we perform a change of basis and a shift,

$$d_{11}(u^{(1)}) = \widehat{d}_{11}(u), \quad d_{12}(u^{(1)}) = \widehat{d}_{12}(u), \quad d_{21}(u^{(1)}) = \widehat{d}_{21}(u), \quad (5.6)$$

$$d_{22}(u^{(1)}) = \widehat{d}_{22}(u) + \psi_1(u^{(1)}) \widehat{d}_{11}(u). \quad (5.7)$$

The function $\psi(u)$ is chosen in such a way that it leads to symmetric exchange relations:

$$\widehat{d}_{12}(u) \widehat{d}_{12}(v) = \begin{cases} \widehat{d}_{12}(v) \widehat{d}_{12}(u), & \text{for } \mathfrak{D}_{2|0}, \mathfrak{D}_{0|2} \\ \mathfrak{h}(u, v) \widehat{d}_{12}(v) \widehat{d}_{12}(u), & \text{for } \mathfrak{D}_{1|1} \end{cases}, \quad (5.8)$$

$$\widehat{d}_{11}(u) \widehat{d}_{12}(v) = \mathfrak{f}_1(u, v) \widehat{d}_{12}(v) \widehat{d}_{11}(u) + \mathfrak{g}_1(u, v) \widehat{d}_{12}(u) \widehat{d}_{11}(v) + \mathfrak{h}_1(u, v) \widehat{d}_{12}(u) \widehat{d}_{22}(v), \quad (5.9)$$

$$\widehat{d}_{22}(u) \widehat{d}_{12}(v) = \widetilde{\mathfrak{f}}_2(u, v) \widehat{d}_{12}(v) \widehat{d}_{22}(u) + \widetilde{\mathfrak{g}}_2(u, v) \widehat{d}_{12}(u) \widehat{d}_{22}(v) + \widetilde{\mathfrak{h}}_2(u, v) \widehat{d}_{12}(v) \widehat{d}_{11}(u). \quad (5.10)$$

The explicit form of the functions appearing above is given in appendix B. In the new basis, Ω is still a pseudo-vacuum:

$$\widehat{d}_{11}(u) \Omega = \widehat{\Lambda}_1(u) \Omega = \Lambda_1(u^{(1)}) \Omega, \quad \widehat{d}_{21}(u) \Omega = 0, \quad (5.11)$$

$$\widehat{d}_{22}(u) \Omega = \widehat{\Lambda}_2(u) \Omega = (\Lambda_2(u^{(1)}) - \psi_1(u^{(1)}) \Lambda_1(u^{(1)})) \Omega. \quad (5.12)$$

and we can use the algebraic Bethe ansatz as in the closed chain case. The transfer matrix rewrites:

$$d(u^{(1)}) = \left((-1)^{[1]} m_1 k(u^{(1)}) + (-1)^{[2]} m_2 \psi_1(u^{(1)}) \right) \widehat{d}_{11}(u) + (-1)^{[2]} m_2 \widehat{d}_{22}(u) \equiv \widehat{d}(u) \quad (5.13)$$

Applying M creation operators $\widehat{d}_{12}(u_j)$ on the pseudo vacuum we generate a Bethe vector:

$$\Phi(\{u\}) = \widehat{d}_{12}(u_1) \dots \widehat{d}_{12}(u_M) \Omega. \quad (5.14)$$

Demanding $\Phi(\{u\})$ to be an eigenvector of $\widehat{d}(u)$ leads to a set of algebraic relations on the parameters $u_1, \dots, u_M$, the so-called Bethe equations. The relation (5.8) between creation operators proves the invariance (up to a function for $\mathfrak{D}_{1|1}$) of the Bethe vector under the reordering of creation operators. This condition is useful to compute the unwanted terms from the action of $\widehat{d}(u)$ on $\Phi(\{u\})$. We compute the action of $\widehat{d}_{11}(u)$ on $\Phi(\{u\})$,

$$\begin{aligned} \widehat{d}_{11}(u) \Phi(\{u\}) &= \prod_{k=1}^M \mathfrak{f}_1(u, u_k) \widehat{\Lambda}_1(u) \Phi(\{u\}) \\ &+ \sum_{k=1}^M \left(M_k(u, \{u\}) \widehat{\Lambda}_1(u_k) + N_k(u, \{u\}) \widehat{\Lambda}_2(u_k) \right) \Phi_k(u, \{u\}), \\ \Phi_k(u, \{u\}) &= \widehat{d}_{12}(u_1) \dots \widehat{d}_{12}(u_k \rightarrow u) \dots \widehat{d}_{12}(u_M) \Omega. \end{aligned} \quad (5.15)$$

where the notation $\widehat{d}_{12}(u_k \rightarrow u)$ is used to indicate the position of $\widehat{d}_{12}(u)$ in the ordered product. The form of $M_1(u; \{u\})$ and $N_1(u; \{u\})$ is easily computed. The other polynomials $M_k(u; \{u\})$ and $N_k(u; \{u\})$ are then computed using the commutation relation between the $\widehat{d}_{12}(u)$ operators and putting $\widehat{d}_{12}(u_k)$ on the left. We get:

$$M_k(u, \{u\}) = \mathfrak{g}_1(u, u_k) \prod_{i \neq k}^M \mathfrak{f}_1(u_k, u_i) \quad \text{and} \quad N_k(u, \{u\}) = \mathfrak{h}_1(u, u_k) \prod_{i \neq k}^M \widetilde{\mathfrak{f}}_2(u_k, u_i).$$

Similarly, we compute the action of $\widehat{d}_{22}(u)$ on $\Phi(\{u\})$,

$$\begin{aligned} \widehat{d}_{22}(u) \Phi(\{u\}) &= \left(\prod_{k=1}^M \widetilde{\mathfrak{f}}_2(u, u_k) \right) \widehat{\Lambda}_2(u) \Phi(\{u\}) + \\ &+ \sum_{k=1}^M \left(O_k(u, \{u\}) \widehat{\Lambda}_2(u_k) + P_k(u, \{u\}) \widehat{\Lambda}_1(u_k) \right) \Phi_k(u, \{u\}), \end{aligned} \quad (5.16)$$

$$O_k(u, \{u\}) = \widetilde{\mathfrak{g}}_2(u, u_k) \prod_{i \neq k}^M \widetilde{\mathfrak{f}}_1(u_k, u_i) \quad \text{and} \quad P_k(u, \{u\}) = \widetilde{\mathfrak{h}}_2(u, u_k) \prod_{i \neq k}^M \mathfrak{f}_1(u_k, u_i). \quad (5.17)$$

Demanding $\Phi(\{u\})$ to be an eigenvector of $\widehat{d}(u)$ corresponds to the cancelling of the so-called ‘unwanted terms’ carried by the vectors $\Phi_k(u, \{u\})$. In this way, we get the Bethe equations:

$$\frac{\widehat{\Lambda}_1(u_k)}{\widehat{\Lambda}_2(u_k)} = \chi_1(u_k) \prod_{i \neq k}^M \frac{\widetilde{\mathfrak{f}}_2(u_k, u_i)}{\mathfrak{f}_1(u_k, u_i)}, \quad k = 1, \dots, M. \quad (5.18)$$

Remark that the r.h.s. depends only on the structure constants of the (super)algebra under consideration, while the l.h.s. encodes the representations entering the spin chain. Then, the eigenvalues of the transfer matrix read:

$$\widehat{d}(u) \Phi(\{u\}) = \widehat{\Lambda}(u; \{u\}) \Phi(\{u\}), \quad (5.19)$$

$$\begin{aligned} \widehat{\Lambda}(u; \{u\}) &= \left((-1)^{[1]} m_1 k(u^{(1)}) + (-1)^{[2]} m_2 \psi_1(u^{(1)}) \right) \widehat{\Lambda}_1(u) \prod_{k=1}^M \mathfrak{f}_1(u, u_k) \\ &+ (-1)^{[2]} m_2 \widehat{\Lambda}_2(u) \prod_{k=1}^M \widetilde{\mathfrak{f}}_2(u, u_k). \end{aligned} \quad (5.20)$$

Note that Bethe equations correspond to the vanishing of the residue of $\Lambda(u; \{u\})$. This is the tool used in analytical Bethe ansatz [12] to obtain Bethe equations, see e.g. [13, 15].

6 Nested Bethe ansatz

6.1 Preliminaries

The method, called the Nested Bethe Ansatz (NBA), consists in a recurrent application of the ABA to express higher rank solutions using the lower ones. It has been introduced in [6] for

the periodic case. The same method can be used for the boundary case. In this way, we can compute the eigenvalues, eigenvectors and Bethe equations of the $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ model from the ones of $\mathfrak{D}_2$ or $\mathfrak{D}_{1|1}$ model. Although we are in a (tensor product of) representation(s) of $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$, we will loosely keep writing $d_{ij}(u)$ the representation of the operators $d_{ij}(u)$, assuming that the reader will understand that when $d_{ij}(u)$ applies to the highest weight Ω , it is in fact its (matricial) representation that is used. Another way to understand this method in an algebraic way is to work in the coset of the $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ algebra by the left ideal $\mathcal{I}_{\mathfrak{m}+\mathfrak{n}}$.

We consider now the open case with general diagonal boundary condition (3.9) for $K^-(u)$, and $K^+(u)$ of the form:

$$K^+(u) = M K(u), \quad \text{with} \quad K(u) = \begin{cases} \mathbb{I} & \text{for } a_+ = 0 \\ \text{diag}(k(u), 1, \dots, 1) & \text{for } a_+ = 1 \end{cases},$$

where the function $k(u)$ is defined in (5.4). The matrix $K^+(u) = M K(u)$ is the only solution we can use to perform the NBA up to the end (see remarks 6.1 and 6.2 below). We decompose the monodromy matrix in the following form (in the $\text{End}(\mathbb{C}^{\mathfrak{m}+\mathfrak{n}})$ auxiliary space),

$$D(u) = \begin{pmatrix} d_{11}(u) & B^{(1)}(u) \\ C^{(1)}(u) & D^{(2)}(u) \end{pmatrix}, \quad (6.1)$$

where $B^{(1)}(u)$ (resp. $C^{(1)}(u)$) is a row (resp. column) vector of $\mathbb{C}^{\mathfrak{m}+\mathfrak{n}-1}$, and $D^{(2)}(u)$ is a matrix of $\text{End}(\mathbb{C}^{\mathfrak{m}+\mathfrak{n}-1})$.

Then, $D^{(2)}(u)$ is itself decomposed in the same way, and more generally, for a given k in $\{1, \dots, \mathfrak{m} + \mathfrak{n} - 2\}$, we gather the generators $d_{kj}(u)$, (resp. $d_{jk}(u)$) $j = k + 1, \dots, \mathfrak{n} + \mathfrak{m}$, in a row (resp. column) vector of $\mathbb{C}^{\mathfrak{m}+\mathfrak{n}-k}$ and $d_{ij}(u)$, $i, j \geq k$, into a matrix of $\text{End}(\mathbb{C}^{\mathfrak{m}+\mathfrak{n}-k})$:

$$B^{(k)}(u) = \sum_{j=k+1}^{\mathfrak{m}+\mathfrak{n}} e_j^t \otimes d_{kj}(u) \quad \text{and} \quad C^{(k)}(u) = \sum_{j=k+1}^{\mathfrak{m}+\mathfrak{n}} e_j \otimes d_{jk}(u), \quad (6.2)$$

$$D^{(k+1)}(u) = \sum_{i,j=k+1}^{\mathfrak{m}+\mathfrak{n}} E_{ij} \otimes d_{ij}(u), \quad (6.3)$$

$$D^{(k)}(u) = \begin{pmatrix} d_{kk}(u) & B^{(k)}(u) \\ C^{(k)}(u) & D^{(k+1)}(u) \end{pmatrix}. \quad (6.4)$$

We decompose the transfer matrix in the same way:

$$\begin{aligned} d(u) &= d^{(1)}(u) = (-1)^{[1]} m_1 k(u) d_{11}(u) + d^{(2)}(u), \\ d^{(k)}(u) &= \text{str}\left(M_a^{(k)} D^{(k)}(u)\right) = (-1)^{[k]} m_k d_{kk}(u) + d^{(k+1)}(u), \end{aligned} \quad (6.5)$$

$$M^{(k)} = \mathbb{I}^{(k)} M \mathbb{I}^{(k)}. \quad (6.6)$$

At each step of the recursion, we make a transformation of the operator and a shift of the spectral parameter:

$$\begin{aligned} d_{kk}(u^{(k)}) &= \widehat{d}_{kk}(u), \quad B_a^{(k)}(u^{(k)}) = \widehat{B}_a^{(k)}(u), \\ D_a^{(k+1)}(u^{(k)}) &= \widehat{D}_a^{(k+1)}(u) + \psi_k(u^{(k)}) \mathbb{I}_a^{(k+1)} \otimes \widehat{d}_{kk}(u). \end{aligned} \quad (6.7)$$

The commutation relations for these operators remain similar for each k :

$$\widehat{B}_a^{(k)}(u) \widehat{B}_b^{(k)}(v) = (-1)^{[k]} \frac{\mathbf{a}_{k+1}(u^{(k)}, v^{(k)})}{\mathbf{a}_k(u^{(k)}, v^{(k)})} \widehat{B}_b^{(k)}(v) \widehat{B}_a^{(k)}(u) \mathbb{R}_{ba}^{(k+1)}(u, v), \quad (6.8)$$

$$\begin{aligned} \widehat{d}_{kk}(u) \widehat{B}_b^{(k)}(v) &= \mathbf{f}_k(u, v) \widehat{B}_b(v) \widehat{d}_{kk}(u) + \mathbf{g}_k(u, v) \widehat{B}_b^{(k)}(u) \widehat{d}_{kk}(v) + \\ &+ \frac{\mathbf{h}_k(u, v)}{\mathbf{e}_{k+1}(v)} \widehat{B}_b^{(k)}(u) \operatorname{str}_a \left(M_a^{(k+1)} \overline{\mathbb{R}}_{ab}^{(k+1)}(v, v) \widehat{D}_a^{(k+1)}(v) \mathbb{R}_{ba}^{(k+1)}(v, v) \right), \end{aligned} \quad (6.9)$$

$$\begin{aligned} \operatorname{str}_a \left(M_a^{(k+1)} \widehat{D}_a^{(k+1)}(u) \right) \widehat{B}_b^{(k)}(v) &= \widetilde{\mathbf{h}}_{k+1}(u, v) \mathbf{e}_{k+1}(u) \widehat{B}_b^{(k)}(u) \widehat{d}_{kk}(v) \\ &+ \frac{\widetilde{\mathbf{g}}_{k+1}(u, v) \mathbf{e}_{k+1}(u)}{\mathbf{e}_{k+1}(v)} \widehat{B}_b^{(k)}(u) \operatorname{str}_a \left(M_a^{(k+1)} \overline{\mathbb{R}}_{ab}^{(k+1)}(v, v) \widehat{D}_a^{(k+1)}(v) \mathbb{R}_{ba}^{(k+1)}(v, v) \right) + \\ &+ \widetilde{\mathbf{f}}_{k+1}(u, v) \widehat{B}_b^{(k)}(v) \operatorname{str}_a \left(M_a^{(k+1)} \overline{\mathbb{R}}_{ab}^{(k+1)}(u, v) \widehat{D}_a^{(k+1)}(u) \mathbb{R}_{ba}^{(k+1)}(u, v) \right). \end{aligned} \quad (6.10)$$

Remark 6.1 *The commutation relations (6.9) and (6.10) impose the restriction on the $K^+(u)$ matrix. The direct use of the reflection equation leads to a matrix $\overline{\mathbb{R}}_{ab}^{(k+1)}(u, u)$ in (6.9) and (6.10). The change $\overline{\mathbb{R}}_{ab}^{(k+1)}(u, u) \rightarrow \overline{\mathbb{R}}_{ab}^{(k+1)}(v, v)$ in the commutation relation is allowed by equality (B.5) which shows that the dependence in u is a scalar function. If the $K^+(u)$ matrix is not from a **NABA couple**, equation (B.5) cannot be used to get (6.9) and (6.10) in their present form. Without this form, the nesting cannot be performed (see also remark 6.2).*

At each step $k = 1, \dots, \mathbf{m} + \mathbf{n} - 1$ of the nesting, we will introduce a family of Bethe parameters u_{kj} , $j = 1, \dots, M_k$, the number M_k of these parameters being a free integer. The partial unions of these families will be noted as,

$$\{u_\ell\} = \bigcup_{i=1}^{\ell} \{u_{ij}, j = 1, \dots, M_i\}, \quad (6.11)$$

so that the whole family of Bethe parameters is $\{u\} = \{u_{\mathbf{m}+\mathbf{n}-1}\}$.

6.2 First step of the construction

From the definition of the highest weight, $C^{(1)}(u)$ annihilates the pseudo-vacuum Ω and we can use $B^{(1)}(u)$ as a creation operator. However, since $B^{(1)}(u)$ contains only $d_{1j}(u)$ operators, it is clear that we need to act on several vectors to describe the whole representation with highest weight Ω . The NBA spirit is to construct these different vectors as Bethe vectors of an $\mathfrak{D}_{\mathbf{m}-1|\mathbf{n}}$ chain that is related to the chain we start with.

More generally, at each step k corresponding to the decomposition (6.4) of the monodromy matrix and to the transformation of the operator of the corresponding algebra $\mathfrak{D}_{\mathbf{m}-k|\mathbf{n}}$, we use (a suitable refinement of) $B^{(k)}(u)$ as a creation operator acting on a set of (to be defined) vectors. These vectors are constructed as Bethe vectors of a $\mathfrak{D}_{\mathbf{m}-k-1|\mathbf{n}}$ chain.

At the first step of the recursion, the Bethe vectors have the form

$$\Phi(\{u^{(1)}\}) = \widehat{B}_{a_1^1}^{(1)}(u_{11}) \dots \widehat{B}_{a_{M_1}^1}^{(1)}(u_{1M_1}) \widehat{F}_{a_1^1 \dots a_{M_1}^1}^{(1)}(\{u\}) \Omega, \quad (6.12)$$

$$\widehat{F}_{a_1^1 \dots a_{M_1}^1}^{(1)}(\{u\}) \in (\mathbb{C}^{\mathbf{m}-1|\mathbf{n}})^{\otimes M_1} \otimes \mathfrak{D}_{\mathbf{m}-1|\mathbf{n}}, \quad (6.13)$$

where $\widehat{F}_{a_1^1 \dots a_{M_1}^1}^{(1)}(\{u\})$ is built from operators $\widehat{d}_{ij}(u)$, $2 \leq i \leq j \leq \mathbf{m} + \mathbf{n}$ only. Since $\widehat{B}^{(1)}(u)$ belongs to $\mathbb{C}^{\mathbf{m}-1|\mathbf{n}} \otimes \mathfrak{D}_{\mathbf{m}|\mathbf{n}}$, we have introduced in the construction M_1 additional auxiliary spaces (labelled $a_1^1, \dots, a_{M_1}^1$) that are also carried by $\widehat{F}_{a_1^1 \dots a_{M_1}^1}^{(1)}(\{u\})$. These new auxiliary spaces take care of the linear combination one has to do between the different generators $\widehat{d}_{1j}(u)$, $j = 2, \dots, \mathbf{m} + \mathbf{n}$, that enter into the construction.

Since $F_{a_1 \dots a_{M_1}}^{(1)}(\{u\})$ is built up from operators $\widehat{d}_{ij}(u)$, $2 \leq i \leq j \leq \mathbf{m} + \mathbf{n}$, it obeys the relation (proven in a more general context in theorem 3.1):

$$\widehat{d}_{11}(u) \widehat{F}_{a_1 \dots a_{M_1}}^{(1)}(\{u\}) \Omega = \widehat{\Lambda}_1(u) \widehat{F}_{a_1 \dots a_{M_1}}^{(1)}(\{u\}) \Omega. \quad (6.14)$$

The transfer matrix is decomposed into

$$d(u^{(1)}) = \widetilde{\mathbf{m}}_1(u) \widehat{d}_{11}(u) + \widehat{d}^{(2)}(u) \quad \text{with} \quad \widehat{d}^{(2)}(u) = \text{str}_a \left(M_a^{(2)} \widehat{D}_a^{(2)}(u) \right) \quad (6.15)$$

$$\widetilde{\mathbf{m}}_1(u) = (-1)^{[1]} m_1 k(u) + \text{str}(M^{(2)}) \psi_1(u^{(1)}). \quad (6.16)$$

The action of $\widehat{d}_{11}(u)$ on $\Phi(\{u^{(1)}\})$ takes the form

$$\begin{aligned} \widehat{d}_{11}(u) \Phi(\{u\}) &= \widehat{\Lambda}_1(u) \prod_{i=1}^{M_1} \mathfrak{f}_1(u, u_{1i}) \Phi(\{u\}) + \sum_{j=1}^{M_1} M_j(u; \{u_1\}) \widehat{\Lambda}_1(u_{1j}) \widehat{\Phi}_j(u, \{u\}) \\ &+ \sum_{j=1}^{M_1} N_j(u; \{u_1\}) \widehat{B}_{a_1}^{(1)}(u_{11}) \dots \widehat{B}_{a_j}^{(1)}(u) \dots \widehat{B}_{a_{M_1}}^{(1)}(u_{1M_1}) \widetilde{d}^{(2)}(u_{1j}; \{u_1\}) \widehat{F}_{a_1 \dots a_{M_1}}^{(1)}(\{u\}) \Omega, \end{aligned} \quad (6.17)$$

with

$$M_j(u; \{u_1\}) = \mathfrak{g}_1(u, u_{1j}) \prod_{i \neq j}^{M_1} \mathfrak{f}_1(u_{1i}, u_{1j}) \quad \text{and} \quad N_j(u; \{u\}) = \frac{\mathfrak{h}_1(u, u_{1j})}{\mathfrak{e}_2(u_{1j})} \prod_{i \neq j}^{M_1} \widetilde{\mathfrak{f}}_2(u_{1i}, u_{1j}).$$

The action of $\widehat{d}^{(2)}(u)$ on $\Phi(\{u\})$ takes the form

$$\begin{aligned} \widehat{d}^{(2)}(u) \Phi(\{u^{(1)}\}) &= \prod_{j=1}^{M_1} \widetilde{\mathfrak{f}}_2(u, u_{1j}) B_{a_1}^{(1)}(u_{11}) \dots \widehat{B}_{a_{M_1}}^{(1)}(u_{1M_1}) \widetilde{d}^{(2)}(u; \{u_1\}) \widehat{F}_{a_1 \dots a_{M_1}}^{(1)}(\{u\}) \Omega \\ &+ \sum_{j=1}^{M_1} P_j(u; \{u_1\}) \widehat{B}_{a_1}^{(1)}(u_{11}) \dots B_{a_j}^{(1)}(u) \dots \widehat{B}_{a_{M_1}}^{(1)}(u_{1M_1}) \widetilde{d}^{(2)}(u_{1j}; \{u_1\}) \widehat{F}_{a_1 \dots a_{M_1}}^{(1)}(\{u\}) \Omega \\ &+ \sum_{j=1}^{M_1} Q_j(u; \{u_1\}) \widehat{\Lambda}_1(u_{1j}) \Phi_j(\{u\}), \end{aligned} \quad (6.18)$$

with

$$P_j(u; \{u_1\}) = \frac{\widetilde{\mathfrak{g}}_2(u, u_{1i}) \mathfrak{e}_2(u)}{\mathfrak{e}_2(u_{1i})} \prod_{i \neq j}^{M_1} \widetilde{\mathfrak{f}}_2(u_{1j}, u_{1i}) \quad ; \quad Q_j(u; \{u_1\}) = \widetilde{\mathfrak{h}}_2(u, u_{1i}) \mathfrak{e}_2(u) \prod_{i \neq j}^{M_1} \mathfrak{f}_1(u_{1j}, u_{1i}).$$

where $\Phi_j(\{u\})$ is deduced from $\Phi(\{u\})$ by the change $u_{1j} \rightarrow u$. These expressions are computed as it has been done in section 5: $N_1(u; \{u_1\})$, $M_1(u; \{u_1\})$, $P_1(u; \{u_1\})$ and $Q_1(u; \{u_1\})$ are easy to compute; the other terms are obtained through a reordering of the operators $\widehat{B}^{(1)}(u_{1j})$, using the reordering lemma 6.1 and the Yang-Baxter equation. We also used the notation:

$$\widetilde{d}^{(2)}(u; \{u_1\}) = \text{str}_a \left(M_a^{(2)} \prod_{j=1}^{\overrightarrow{M_1}} \overline{\mathbb{R}}_{a,a_j}^{(2)}(u, u_{1j}) \widehat{D}_a^{(2)}(u) \prod_{j=1}^{\overleftarrow{M_1}} \mathbb{R}_{a_j,a}^{(2)}(u, u_{1j}) \right). \quad (6.19)$$

Remark 6.2 *The fact that the wanted and unwanted terms contain the same operator $\widetilde{d}^{(2)}$ (but at different values u and u_{1j}) allows to continue the nesting. In this way, the diagonalisation of this operator allows at the same time to compute the eigenvalue and to show that the unwanted terms cancel (when the Bethe ansatz equations are obeyed). The apparition of this operator in the unwanted terms is directly related to the present form of the commutation relations (6.9) and (6.10), see remark 6.1. Hence the need of a NABA couple to perform the nesting.*

As already mentionned, the calculation makes appear a new transfer matrix $\widetilde{d}^{(2)}(u; \{u_1\})$ corresponding to a $\mathfrak{D}_{m-1|n}$ chain with $L + M_1$ sites, the M_1 additional sites corresponding to fundamental representations of $\mathfrak{D}_{m-1|n}$. This interpretation is supported by theorem 3.1 which ensure that $\widetilde{D}_a^{(2)}(u; \{u_1\})$ generates $\mathfrak{D}_{m-1|n}$, and that $\widetilde{d}^{(2)}(u; \{u_1\})$ is indeed the transfer matrix of an integrable spin chain. Then, if we assume that $F_{a_1 \dots a_{M_1}}^{(1)}(\{u\}) \Omega$ is an eigenvector of this new transfer matrix,

$$\widetilde{d}^{(2)}(u; \{u_1\}) \widehat{F}_{a_1 \dots a_{M_1}}^{(1)}(\{u\}) \Omega = \widetilde{\Gamma}^{(2)}(u) \widehat{F}_{a_1 \dots a_{M_1}}^{(1)}(\{u\}) \Omega, \quad (6.20)$$

we deduce:

$$\begin{aligned} \widehat{d}_{11}(u) \Phi(\{u\}) &= \widehat{\Lambda}_1(u) \prod_{i=1}^{M_1} \mathfrak{f}_1(u, u_{1i}) \Phi(\{u\}) \\ &+ \sum_{j=1}^{M_1} \left(M_j(u; \{u_1\}) \widehat{\Lambda}_1(u_{1j}) + N_j(u; \{u_1\}) \widetilde{\Gamma}^{(2)}(u_{1j}) \right) \Phi_j(\{u\}), \end{aligned} \quad (6.21)$$

$$\begin{aligned} \widehat{d}^{(2)}(u) \Phi(\{u\}) &= \widetilde{\Gamma}^{(2)}(u) \prod_{j=1}^{M_1} \widetilde{\mathfrak{f}}_2(u, u_{1j}) \Phi(\{u\}) \\ &+ \sum_{j=1}^{M_1} \left(P_j(u; \{u_1\}) \widetilde{\Gamma}^{(2)}(u_{1j}) + Q_j(u; \{u_1\}) \widehat{\Lambda}_1(u_{1j}) \right) \Phi_j(\{u\}). \end{aligned} \quad (6.22)$$

Gathering these relations together, we get a first expression of the action of $d(u)$ on $\Phi(\{u\})$. When we cancel in this expression the unwanted terms (carried by $\Phi_j(\{u\})$), we get the first system of Bethe equations:

$$\frac{\widehat{\Lambda}_1(u_{1j})}{\widetilde{\Gamma}^{(2)}(u_{1j})} = \frac{\chi_1(u_{1j})}{\mathfrak{e}_2(u_{1j})} \prod_{i \neq j}^{M_1} \frac{\widetilde{\mathfrak{f}}_2(u_{1j}, u_{1i})}{\mathfrak{f}_1(u_{1j}, u_{1i})}. \quad (6.23)$$

We also get a first expression of the transfer matrix eigenvalue:

$$\widehat{d}^{(1)}(u) \Phi(\{u\}) = \left(\widetilde{m}_1(u) \widehat{\Lambda}_1(u) \prod_{j=1}^{M_1} \widehat{f}_1(u, u_{1j}) + \widetilde{\Gamma}^{(2)}(u) \prod_{j=1}^{M_1} \widetilde{f}_1(u, u_{1j}) \right) \Phi(\{u\}). \quad (6.24)$$

In the above relations, everything is known *but* the eigenvalue $\widetilde{\Gamma}^{(2)}(u)$, introduced in (6.20), and the explicit form of $\widehat{F}_{a_1^1 \dots a_{M_1}^1}^{(1)}(\{u\})$ ensuring that (6.20) is indeed satisfied.

Thus, at the end of this first recursion step, we have ‘reduced’ the problem of computing an eigenvector $\Phi(\{u\})$ for the transfer matrix $d(u)$ of a $\mathfrak{D}_{m|n}$ chain with L sites to the problem of computing an eigenvector $\Phi^{(1)}(\{u\}) = \widehat{F}_{a_1^1 \dots a_{M_1}^1}^{(1)}(\{u\}) \Omega$ for the transfer matrix $\widetilde{d}^{(2)}(u; \{u_1\})$ of a $\mathfrak{D}_{m-1|n}$ chain with $L + M_1$ sites.

Remark 6.3 (Change of notation) *To avoid too complicated notation for the second step, we need to slightly change the notation at the end of the first step. First, we rename the hatted operators $\widehat{X}(u) \rightarrow X(u)$, although they still have the spectral parameter shift and the operator transformation coming from the first step. Secondly, we omit the tilda on operators, $\widetilde{X}(u) \rightarrow X(u)$, keeping in mind that the new operator $X(u)$ have M_1 sites more than the one of the previous step. In this way, we will be able to re-use the ‘hatted’ and ‘tilded’ notations for the transformations used in the second step.*

This will be the general approach at each step: at the end of step k , we will perform a change of notation, suppressing the hats and tildas on operators, to use them again in step $k + 1$.

It remains to single out the highest weights corresponding to the fundamental representations carried by the new sites. This is done in the following way

$$\begin{aligned} \Phi^{(1)}(\{u\}) &= F_{a_1^1 \dots a_{M_1}^1}^{(1)}(\{u\}) \Omega, \\ \Phi^{(1)}(\{u\}) &= B_{a_1^2}^{(2)}(u_{21}; \{u_1\}) \dots B_{a_{M_2}^2}^{(2)}(u_{2M_2}; \{u_1\}) F_{a_1^2 \dots a_{M_2}^2}^{(2)}(\{u\}) \Omega^{(2)}, \end{aligned} \quad (6.25)$$

$$\Omega^{(2)} = (e_1^{(1)})^{\otimes M_1} \otimes \Omega, \quad (6.26)$$

where $e_1^{(1)} = (1, 0, \dots, 0)^t \in \mathbb{C}^{m-1|n}$ and $F_{a_1^2 \dots a_{M_2}^2}^{(2)}(\{u\})$ is built on operators $d_{ij}(u^{(2)}; \{u_1\})$, with $j \geq i > 2$. The operators $B^{(2)}(u; \{u_1\})$ play the role, for the $\mathfrak{D}_{m-1|n}$ chain of length $L + M_1$, of the operators $B^{(1)}(u)$ for the $\mathfrak{D}_{m|n}$ chain of length L . Explicitly, they are obtained from the decomposition (6.4) of the monodromy matrix.

6.3 General step

More generally, the step k starts with the problem

$$d^{(k)}(u; \{u_{k-1}\}) \Phi^{(k-1)}(\{u\}) = \Gamma^{(k)}(u) \Phi^{(k-1)}(\{u\}), \quad (6.27)$$

where $d^{(k)}(u; \{u_{k-1}\}) = \text{str} \left(M^{(k)} D^{(k)}(u; \{u_{k-1}\}) \right)$ is the transfer matrix of a $\mathfrak{D}_{m-k-1|n}$ spin chain of length $L + \sum_{j=1}^{k-1} M_j$ (obtained from the previous step). We recall that hats and tildas

have been suppressed, according to remark 6.3, including for the function $\Gamma^{(k)}(u)$. We define

$$\Phi^{(k-1)}(\{u\}) = F_{a_1^{k-1} \dots a_{M_{k-1}}^{k-1}}^{(k-1)}(\{u\}) \Omega^{(k-1)} = \mathbb{B}^{(k)}(\{u_k\}) F_{a_1^k \dots a_{M_k}^k}^{(k)}(\{u\}) \Omega^{(k)}, \quad (6.28)$$

$$\Omega^{(k)} = (e_1^{(k-1)})^{\otimes M_{k-1}} \otimes \Omega^{(k-1)}, \quad (6.29)$$

with $e_1^{(k)} = (1, 0, \dots, 0)^t \in \mathbb{C}^{m-k|n}$. We have introduced

$$\mathbb{B}^{(k)}(\{u_k\}) = B_{a_1^k}^{(k)}(u_{k1}; \{u_{k-1}\}) \dots B_{a_{M_k}^k}^{(k)}(u_{kM_k}; \{u_{k-1}\}), \quad (6.30)$$

where the operators are extracted from the monodromy matrix, see equation (6.4).

Remark 6.4 In (6.30), we have indicated only the auxiliary spaces a_j^k , $j = 1, \dots, M_k$. In fact, since $D^{(k)}(u; \{u_{k-1}\})$ is viewed as the monodromy matrix of a spin chain of length $L + \sum_{j=1}^{k-1} M_j$, the other spaces a_j^ℓ , $j = 1, \dots, M_\ell$, $\ell < k$, are now quantum spaces. Thus, they do not appear explicitly in $D^{(k)}$, as the sites of the original spin chain, but obviously this monodromy matrix (and its components) does depend on all these spaces.

We extract from $d^{(k)}(u; \{u_{k-1}\})$ the component $d_{kk}(u; \{u_{k-1}\})$:

$$d^{(k)}(u; \{u_{k-1}\}) = (-1)^{[k]} m_k d_{kk}(u; \{u_{k-1}\}) + \text{str} \left(M^{(k+1)} D^{(k+1)}(u; \{u_{k-1}\}) \right). \quad (6.31)$$

Now we must transform the operator:

$$\begin{aligned} d_{kk}(u^{(k)}) &= \widehat{d}_{kk}(u), & B_a^{(k)}(u^{(k)}) &= \widehat{B}_a^{(k)}(u), \\ D^{(k+1)}(u^{(k)}) &= \widehat{D}^{(k+1)}(u) + \psi_k(u^{(k)}) \widehat{d}_{kk}(u) \mathbb{I}^{(k)}. \end{aligned} \quad (6.32)$$

The transfer matrix $d^{(k)}(u^{(k)}; \{u_{k-1}\})$ is rewritten as

$$\widehat{d}^{(k)}(u; \{u_{k-1}\}) = \widetilde{\mathbf{m}}_k(u) \widehat{d}_{kk}(u; \{u_{k-1}\}) + \text{str} \left(M^{(k+1)} \widehat{D}^{(k+1)}(u; \{u_{k-1}\}) \right), \quad (6.33)$$

$$\widetilde{\mathbf{m}}_k(u) = (-1)^{[k]} m_k + \text{str}(M^{(k)}) \psi_k(u^{(k)}), \quad (6.34)$$

and the Bethe vector:

$$\Phi^{(k-1)}(\{u\}) = \widehat{\mathbb{B}}^{(k)}(\{u_k\}) \widehat{F}_{a_1^k \dots a_{M_k}^k}^{(k)}(\{u\}) \Omega^{(k)}, \quad (6.35)$$

$$\widehat{\mathbb{B}}^{(k)}(\{u_k\}) = \widehat{B}_{a_1^k}^{(k)}(u_{k1}; \{u_{k-1}\}) \dots \widehat{B}_{a_{M_k}^k}^{(k)}(u_{kM_k}; \{u_{k-1}\}). \quad (6.36)$$

Now we can compute the action of the transfer matrix on this vector. We first commute $\widehat{d}_{kk}(u; \{u_{k-1}\})$ and $\widehat{d}^{(k+1)}(u; \{u_{k-1}\}) = \text{str} \left(M^{(k)} \widehat{D}^{(k+1)}(u; \{u_{k-1}\}) \right)$ with the operator $\widehat{\mathbb{B}}^{(k)}(\{u_k\})$:

$$\begin{aligned} \widehat{d}_{kk}(u; \{u_{k-1}\}) \Phi^{(k-1)}(\{u^{(k)}\}) &= \prod_{j=1}^{M_k} \mathbf{f}_k(u, u_{kj}) \widehat{\mathbb{B}}^{(k)}(\{u_k\}) \widehat{d}_{kk}(u; \{u_{k-1}\}) \Phi^{(k)}(\{u\}) \\ &+ \sum_{j=1}^{M_k} M_j(u; \{u_{k-1}\}) \widehat{\mathbb{B}}_j^{(k)}(u; \{u_k\}) \widehat{d}_{kk}(u_{kj}; \{u_{k-1}\}) \Phi^{(k)}(\{u\}) \\ &+ \sum_{j=1}^{M_k} N_j(u; \{u_{k-1}\}) \widehat{\mathbb{B}}_j^{(k)}(u; \{u_k\}) \widehat{d}^{(k+1)}(u_{kj}; \{u_k\}) \Phi^{(k)}(\{u\}), \end{aligned} \quad (6.37)$$

$$\begin{aligned}
\widehat{d}^{(k+1)}(u; \{u_{k-1}\}) \Phi^{(k-1)}(\{u\}) &= \prod_{j=1}^{M_k} \widetilde{f}_{k+1}(u, u_{kj}) \widehat{\mathbb{B}}^{(k)}(\{u_k\}) \widetilde{d}^{(k+1)}(u; \{u_k\}) \Phi^{(k)}(\{u\}) \\
&+ \sum_{j=1}^{M_k} P_j(u; \{u_{k-1}\}) \widehat{\mathbb{B}}_j^{(k)}(u; \{u_k\}) \widetilde{d}^{(k+1)}(u_{kj}; \{u_k\}) \Phi^{(k)}(\{u\}) \\
&+ \sum_{j=1}^{M_k} Q_j(u; \{u_{k-1}\}) \widehat{\mathbb{B}}_j^{(k)}(u; \{u_k\}) \widehat{d}_{kk}(u_{kj}; \{u_{k-1}\}) \Phi^{(k)}(\{u\}), \tag{6.38}
\end{aligned}$$

where we have introduced:

$$\begin{aligned}
\Phi^{(k)}(\{u\}) &= \widehat{F}_{a_1^k \dots a_{M_k}^k}^{(k)}(\{u\}) \Omega^{(k)}, \\
\widetilde{d}^{(k+1)}(u; \{u_k\}) &= \text{str}_a \left(M_a^{(k+1)} \left(\prod_{j=1}^{\overrightarrow{M_k}} \overline{\mathbb{R}}_{aa_j^k}^{(k+1)}(u, u_{kj}) \right) \widehat{D}_a^{(k+1)}(u; \{u_{k-1}\}) \left(\prod_{j=1}^{\overleftarrow{M_k}} \mathbb{R}_{a_j^k a}^{(k+1)}(u, u_{kj}) \right) \right).
\end{aligned}$$

The functions M_j , N_j , P_j and Q_j are the same as in the first step (section 6.2) but with indices $1 \rightarrow k$ on functions and Bethe roots. We use the following reordering lemma:

Lemma 6.1 *For each $k = 1, \dots, \mathfrak{m} + \mathfrak{n} - 1$ and $j = 1, \dots, M_k$, we have*

$$\begin{aligned}
\widehat{\mathbb{B}}^{(k)}(\{u_k\}) &= \widehat{B}_j^{(k)}(u_{kj}) \widehat{B}_1^{(k)}(u_{k1}) \dots \widehat{B}_{j-1}^{(k)}(u_{k,j-1}) \widehat{B}_{j+1}^{(k)}(u_{k,j+1}) \dots \widehat{B}_{M_k}^{(k)}(u_{kM_k}) \\
&\times \prod_{i=1}^{\overrightarrow{j-1}} (-1)^{[j]} \frac{\mathfrak{a}_{j+1}(u_{ki}^{(k)}, u_{kj}^{(k)})}{\mathfrak{a}_j(u_{ki}^{(k)}, u_{kj}^{(k)})} \mathbb{R}_{ji}^{(k+1)}(u_{ki}, u_{kj}) \tag{6.39}
\end{aligned}$$

where the dependence in $\{u_{k-1}\}$ has been omitted in $\widehat{B}_p^{(k)}$.

Proof: Direct calculation using the commutation relations (6.8)-(6.10). ■

We now compute the action of $\widehat{d}_{kk}(u; \{u_{k-1}\})$ and $\widetilde{d}^{(k+1)}(u; \{u_k\})$ on $\widehat{F}^{(k)}(\{u\}) \Omega^{(k)}$. These actions follow from the theorem 3.1. For $\widehat{d}_{kk}(u; \{u_{k-1}\})$ we have:

$$\widehat{d}_{kk}(u; \{u_{k-1}\}) \widehat{F}^{(k)}(\{u\}) \Omega^{(k)} = \widehat{\Lambda}_k(u; \{u_{k-1}\}) \widehat{F}^{(k)}(\{u\}) \Omega^{(k)}. \tag{6.40}$$

It remains to do the same for $\widetilde{d}^{(k+1)}(u; \{u_k\})$. It corresponds to a new monodromy matrix

$$\widetilde{D}_a^{(k+1)}(u; \{u_k\}) = \prod_{j=1}^{\overrightarrow{M_k}} \overline{\mathbb{R}}_{aa_j^k}^{(k+1)}(u, u_{kj}) \widehat{D}_a^{(k+1)}(u; \{u_{k-1}\}) \prod_{j=1}^{\overleftarrow{M_k}} \mathbb{R}_{a_j^k a}^{(k+1)}(u, u_{kj}). \tag{6.41}$$

It also satisfies the reflexion equation, see the theorem 3.1, so that the problem is integrable, and defines a $\mathfrak{D}_{\mathfrak{m}-k|\mathfrak{n}}$ spin chain, with $L + \sum_{j=1}^k M_j$ sites.

We get a new eigenvalue problem:

$$\widetilde{d}^{(k+1)}(u; \{u_k\}) \Phi^{(k)}(\{u\}) = \widetilde{\Gamma}^{(k+1)}(u) \Phi^{(k)}(\{u\}). \tag{6.42}$$

Assuming the form (6.42), we can show, following the same lines as in the first step, that $\Phi^{(k-1)}(\{u\})$ is a transfer matrix eigenvector provided the k^{th} system of Bethe equations,

$$\frac{\widehat{\Lambda}_k(u_{kj}; \{u_{k-1}\})}{\widehat{\Gamma}^{(k+1)}(u_{kj}; \{u_k\})} = \frac{\chi_k(u_{kj})}{\mathfrak{e}_{k+1}(u_{kj})} \prod_{i \neq j}^{M_k} \frac{\widetilde{f}_{k+1}(u_{kj}, u_{ki})}{f_k(u_{kj}, u_{ki})}, \quad (6.43)$$

is obeyed. We also get an expression for $\widehat{\Gamma}^{(k)}(u)$, the eigenvalue of $\widehat{d}^{(k)}(u)$:

$$\widehat{\Gamma}^{(k)}(u) = \widetilde{m}_k(u) \prod_{j=1}^{M_k} f_k(u, u_{kj}) \widehat{\Lambda}_k(u; \{u_{k-1}\}) + \prod_{j=1}^{M_k} \widetilde{f}_{k+1}(u, u_{kj}) \widetilde{\Gamma}^{(k+1)}(u; \{u_k\}). \quad (6.44)$$

6.4 End of induction

To end the recursion, we use the $\mathbf{m} + \mathbf{n} = 2$ case and remark that:

$$\widehat{\Gamma}^{(\mathbf{n}+\mathbf{m})}(u) = \widehat{\Lambda}_{\mathbf{n}+\mathbf{m}}(u, \{u_{\mathbf{m}+\mathbf{n}-2}\}). \quad (6.45)$$

Using the shift notation, $u^{(k \dots l)} = (\dots (u^{(k)})^{(k+1)} \dots)^{(l)}$, for $k \leq l$, we deduce from (6.45) that $\widehat{\Gamma}$ is expressed in term of $\widehat{\Lambda}$:

$$\begin{aligned} \widehat{\Gamma}^{(k+1)}(u^{(k+2 \dots \mathbf{n}+\mathbf{m}-1)}) &= \widetilde{m}_{k+1}(u^{(k+2 \dots \mathbf{n}+\mathbf{m}-1)}) \widehat{\Lambda}_{k+1}(u^{(k+2 \dots \mathbf{n}+\mathbf{m}-1)}; \{u_k\}) \mathcal{F}_{k+1}(u) \\ &+ \sum_{\ell=k+2}^{\mathbf{m}+\mathbf{n}-1} \widetilde{m}_\ell(u^{(\ell+1 \dots \mathbf{n}+\mathbf{m}-1)}) \widehat{\Lambda}_\ell(u^{(\ell+1 \dots \mathbf{n}+\mathbf{m}-1)}; \{u_{\ell-1}\}) \mathcal{F}_\ell(u) \prod_{p=k+1}^{\ell-1} \widetilde{\mathcal{F}}_p(u) \\ &+ (-1)^{[\mathbf{m}+\mathbf{n}]} m_{\mathbf{m}+\mathbf{n}} \widehat{\Lambda}_{\mathbf{m}+\mathbf{n}}(u; \{u_{\mathbf{m}+\mathbf{n}-2}\}) \prod_{p=k+1}^{\mathbf{m}+\mathbf{n}-1} \widetilde{\mathcal{F}}_p(u), \end{aligned} \quad (6.46)$$

where we have introduced

$$\mathcal{F}_\ell(u) = \prod_{j=1}^{M_\ell} f_\ell(u^{(\ell+1 \dots \mathbf{n}+\mathbf{m}-1)}, u_{\ell j}), \quad \widetilde{\mathcal{F}}_\ell(u) = \prod_{j=1}^{M_\ell} \widetilde{f}_{\ell+1}(u^{(\ell+1 \dots \mathbf{n}+\mathbf{m}-1)}, u_{\ell j}), \quad \ell \in \{k \dots \mathbf{m} + \mathbf{n} - 1\},$$

with the convention $u^{(k \dots l)} = u$ if $k > l$. It remains to compute the values $\widehat{\Lambda}_k(u; \{u_{k-1}\})$:

Lemma 6.2 *The eigenvalue $\widehat{\Lambda}_k(u; \{u_{k-1}\})$ of $\widehat{d}_{kk}(u; \{u_{k-1}\})$ on $\Omega^{(k-1)}$ is given by,*

$$\widehat{\Lambda}_k(u; \{u_k\}) = \widehat{\Lambda}_k(u) \prod_{\ell=1}^{k-2} \prod_{j=1}^{M_\ell} \frac{1}{\widetilde{f}_{\ell+1}(u^{(\ell+1 \dots k)}, u_{\ell j})} = \widehat{\Lambda}_k(u) \prod_{\ell=1}^{k-2} \frac{1}{\widetilde{\mathcal{F}}_\ell(u^{(k+1 \dots \mathbf{m}+\mathbf{n}-1)})}, \quad k = 1, \dots, \mathbf{m}+\mathbf{n}, \quad (6.47)$$

where we have used $\widehat{d}_{kk}(u) \Omega = \widehat{\Lambda}_k(u) \Omega$ with:

$$\widehat{\Lambda}_k(u) = \Lambda_k(u^{(1 \dots k)}) - \sum_{i=1}^{k-1} q^{2(k-1-i)-4 \sum_{l=i+1}^{k-1} [l]} \psi_i((u^{(k)})^{(i)}) \Lambda_i(u^{(1 \dots k)}) \quad (6.48)$$

$$= \mathcal{K}_k(u^{(1 \dots k)}) \lambda_k(u^{(1 \dots k)}) \lambda'_k(u^{(1 \dots k)}). \quad (6.49)$$

Proof: First we introduce a useful property between coproduct and supertrace:

$$\text{str}_a(\Delta(D_a^{(k)}(u)) = \Delta(\text{str}_a(D_a^{(k)}(u))). \quad (6.50)$$

It is obvious because supertrace and coproduct do not act in the same space. We recall the fundamental representation evaluation map for the $\mathcal{A}_{\mathbf{m}-k+1|\mathbf{n}}$ algebra:

$$\begin{aligned} \mathcal{A}_{\mathbf{m}-k+1|\mathbf{n}} \otimes \text{End}(\mathbb{C}^{\mathbf{m}|\mathbf{n}}) &\rightarrow \text{End}(\mathbb{C}^{\mathbf{m}-k+1|\mathbf{n}}) \otimes \text{End}(\mathbb{C}^{\mathbf{m}|\mathbf{n}}) \\ \pi_a^{(k)} : T^{(k)}(u) &\mapsto \overline{\mathbb{R}}_{12}^{(k)}(u, a) \\ (T^{-1}(\boldsymbol{\iota}(u)))^{(k)} &\mapsto \mathbb{R}_{21}^{(k)}(u, a) \end{aligned} \quad (6.51)$$

We also need the representation of $\mathcal{A}_{\mathbf{m}-k+1|\mathbf{n}}$ induced by the inclusion $\mathcal{A}_{\mathbf{m}-k+1|\mathbf{n}} \hookrightarrow \mathcal{A}_{\mathbf{m}-p+1|\mathbf{n}}$, $p < k$. From the identity

$$T^{(k)}(u) = \mathbb{I}^{(k)}(T^{(p)}(u^{(p+1\dots k)})) \mathbb{I}^{(k)}, \quad (6.52)$$

we can deduce the form of $\pi_v^{(p)}(T^{(k)}(u))$ in the fundamental representation of $\mathcal{A}_{\mathbf{m}-p+1|\mathbf{n}}$:

$$\begin{aligned} (id \otimes \pi_v^{(p)})(T^{(k)}(u)) &= \mathbb{I}_1^{(k)}(id \otimes \pi_v^{(p)})(T^{(p)}(u^{(p+1\dots k)})) \mathbb{I}_1^{(k)} = \mathbb{I}_1^{(k)} \overline{\mathbb{R}}_{12}^{(p)}(u^{(p+1\dots k)}, v) \mathbb{I}_1^{(k)} \\ &= \overline{\mathbb{R}}_{a,b}^{(k,p)}(u^{(p+1\dots k)}, v). \end{aligned} \quad (6.53)$$

The last equality is just the definition of $\mathbb{R}_{a,b}^{(k,p)}(u^{(p+1\dots k)}, v)$, see (2.15).

Hence, using theorem 3.3, we can rewrite the monodromy operator $\tilde{D}_a^{(k+1)}(u; \{u_{k-1}\})$ as:

$$\begin{aligned} \tilde{D}_a^{(k+1)}(u; \{u_{k-1}\}) &= \left(\prod_{j=1}^{\overrightarrow{M_k}} \overline{\mathbb{R}}_{aa_j^k}^{(k+1)}(u, u_{kj}) \right) \hat{D}_a^{(k+1)}(u; \{u_{k-1}\}) \left(\prod_{j=1}^{\overleftarrow{M_k}} \mathbb{R}_{a_j^k a}^{(k+1)}(u, u_{kj}) \right) \\ &= (id \otimes (\pi_{u_{ik}}^{(k+1)})^{\otimes_{i=1}^{M_k}}) \circ \Delta^{M_k}(\hat{D}_a^{(k+1)}(u; \{u_{k-1}\})), \end{aligned} \quad (6.54)$$

while the operator $\tilde{d}_{kk}^{(k)}(u)$ takes the form:

$$\tilde{d}_{kk}^{(k)}(u; \{u_{k-1}\}) = (id \otimes ((\pi_{u_{ip}}^{(p+1)})^{\otimes_{i=1}^{M_p}})^{\otimes_{p=1}^{k-1}}) \circ \Delta^{\sum_{p=1}^k M_p}(\tilde{d}_{kk}^{(k)}(u)). \quad (6.55)$$

Now acting on the highest weight $\Omega^{(k-1)}$ and using lemma 3.2, we find the following result:

$$\tilde{d}_{kk}^{(k)}(u) \Omega^{(k-1)} = (id \otimes ((\pi_{u_{ip}}^{(p+1)})^{\otimes_{i=1}^{M_p}})^{\otimes_{p=1}^k})(t_{kk}^{(k)}(u) t_{kk}'^{(k)}(\boldsymbol{\iota}(u)))^{\otimes_{p=1}^k M_p} \otimes \tilde{d}_{kk}^{(k)}(u) \Omega^{(k-1)}. \quad (6.56)$$

We have from the definition of $\mathbb{R}$ matrices and $\Omega^{(k-1)}$,

$$\begin{aligned} \pi_{u_{ip}}^{(p+1)}(t_{kk}^{(k)}(u) t_{kk}'^{(k)}(\boldsymbol{\iota}(u))) \Omega^{(k-1)} &= \frac{\mathbf{b}(u^{(p+1\dots k)}, u_{ip}) \bar{\mathbf{b}}(u^{(p+1\dots k)}, u_{ip})}{\mathbf{a}_{p+1}(u^{(p+1\dots k)}, u_{ip}) \bar{\mathbf{a}}_{p+1}(u^{(p+1\dots k)}, u_{ip})} \Omega^{(k-1)}, \\ \pi_{u_{ik}}^{(k)}(t_{kk}^{(k)}(u) t_{kk}'^{(k)}(\boldsymbol{\iota}(u))) \Omega^{(k-1)} &= \Omega^{(k-1)}, \end{aligned} \quad (6.57)$$

that leads to the result (6.47). The eigenvalue (6.48) is computed directly from theorem 3.1. To obtain the form (6.49), one uses the equalities (4.38), (4.39) and the identity (B.10). ■

From the expression given in lemma 6.2, one deduces that:

$$\begin{aligned}\widehat{\Gamma}^{(k+1)}(u^{(k+2\dots n+m-1)}) &= \left\{ \widetilde{m}_{k+1}(u^{(k+2\dots n+m-1)}) \widehat{\Lambda}_{k+1}(u^{(k+2\dots n+m-1)}) \mathcal{F}_{k+1}(u) \right. \\ &\quad + \sum_{\ell=k+2}^{m+n-1} \widetilde{m}_{\ell}(u^{(\ell+1\dots n+m-1)}) \widehat{\Lambda}_{\ell}(u^{(\ell+1\dots n+m-1)}) \mathcal{F}_{\ell}(u) \widetilde{\mathcal{F}}_{\ell-1}(u) \\ &\quad \left. + (-1)^{[m+n]} m_{m+n} \widehat{\Lambda}_{m+n}(u^{(n+m-1)}) \widetilde{\mathcal{F}}_{m+n-1}(u) \right\} \frac{1}{\prod_{\ell=1}^{k-1} \widetilde{\mathcal{F}}_{\ell}(u)} \quad (6.58)\end{aligned}$$

Let us note that since $\mathfrak{b}(u, u) = 0$, equation (6.58) implies that:

$$\widehat{\Gamma}^{(k+1)}(u_{\ell j}) = 0, \quad \text{for } j = 1, \dots, M_{\ell}; \ell = 1, \dots, k-1, \quad (6.59)$$

$$\begin{aligned}\widehat{\Gamma}^{(k+1)}(u_{kj}) &= \widetilde{m}_{k+1}(u_{kj}) \widehat{\Lambda}_{k+1}(u_{kj}) \mathcal{F}_{k+1}(u_{kj}^{\overline{(k+2\dots n+m-1)}}) \prod_{\ell=1}^{k-1} \frac{1}{\widetilde{\mathcal{F}}_{\ell+1}(u_{kj}^{\overline{(k+2\dots n+m-1)}})}, \\ &\quad \text{for } j = 1, \dots, M_k. \quad (6.60)\end{aligned}$$

6.5 Final form of Bethe vectors, eigenvalues and equations

Using expressions (6.59), (6.60), and the value of $\widehat{\Lambda}_k(u; \{u^{(k)}\})$ given in lemma 6.2, one can recast the Bethe equations (6.43) in their final form:

$$\begin{aligned}\frac{\widehat{\Lambda}_k(u_{kj})}{\widehat{\Lambda}_{k+1}(u_{kj}^{\overline{(k+1)}})} &= \frac{\widetilde{m}_{k+1}(u_{kj}^{\overline{(k+1)}}) \chi_k(u_{kj})}{\mathfrak{e}_{k+1}(u_{kj})} \prod_{i=1}^{M_{k-1}} \frac{1}{\widetilde{f}_k(u_{kj}^{(k)}, u_{k-1,i})} \prod_{i \neq j}^{M_k} \frac{\widetilde{f}_{k+1}(u_{kj}, u_{ki})}{f_k(u_{kj}, u_{ki})} \prod_{i=1}^{M_{k+1}} f_{k+1}(u_{kj}^{\overline{(k+1)}}, u_{k+1,i}), \\ &\quad j = 1, \dots, M_k, \quad k = 1, \dots, m+n-1, \quad (6.61)\end{aligned}$$

with the convention $M_0 = M_{m+n} = 0$. The number of parameter families is $m+n-1$. We checked that using the weights (4.3), (4.5) and the functions given in appendix B, one reproduces the BAEs already computed, in e.g. [6, 12, 17], and also the general forms given in [13–15]. In particular, for the fundamental weight $\mu = (1, 0, \dots, 0)$, we recovers the BAEs for a spin chain with fundamental representations. For instance, for the case of $\widehat{\mathcal{U}}_q(2|2)$, which may be of some relevance in the context of AdS/CFT correspondence, one gets (for L sites with evaluation parameter b_i , $a_+ = 1$ and $a_- = 2$):

$$\begin{aligned}&\frac{c_+ u_{1i} q^{\frac{1}{2}} - \frac{q^{-\frac{1}{2}}}{c_+ u_{1i}}}{\frac{u_{1i}}{c_+} q^{\frac{1}{2}} - \frac{c_+}{u_{1i}} q^{-\frac{1}{2}}} \prod_{i=1}^L \frac{(\frac{u_{1i}}{b_i} q^{\frac{1}{2}} - \frac{b_i}{u_{1i}} q^{-\frac{1}{2}})(u_{1i} b_i q^{\frac{1}{2}} - \frac{q^{-\frac{1}{2}}}{b_i u_{1i}})}{(\frac{u_{1i}}{b_i} q^{-\frac{1}{2}} - \frac{b_i}{u_{1i}} q^{\frac{1}{2}})(u_{1i} b_i q^{-\frac{1}{2}} - \frac{q^{\frac{1}{2}}}{b_i u_{1i}})} \\ &= \prod_{j \neq i}^{M_1} \frac{(\frac{u_{1j}}{u_{1i}} q - \frac{u_{1j}}{u_{1i}} q^{-1})(u_{1i} u_{1j} q - \frac{q^{-1}}{u_{1i} u_{1j}})}{(\frac{u_{1j}}{u_{1i}} q - \frac{u_{1j}}{u_{1i}} q^{-1})(u_{1i} u_{1j} q^{-1} - \frac{q}{u_{1i} u_{1j}})} \prod_{j=1}^{M_2} \frac{(\frac{u_{2j}}{u_{1i}} q^{-\frac{1}{2}} - \frac{u_{1i}}{u_{2j}} q^{\frac{1}{2}})(u_{2j} u_{1i} q^{-\frac{1}{2}} - \frac{q^{\frac{1}{2}}}{u_{2j} u_{1i}})}{(\frac{u_{2j}}{u_{1i}} q^{-\frac{1}{2}} - \frac{u_{2j}}{u_{1i}} q^{\frac{1}{2}})(u_{2j} u_{1i} q^{\frac{1}{2}} - \frac{q^{-\frac{1}{2}}}{u_{2j} u_{1i}})}, \\ &\quad i = 1, \dots, M_1\end{aligned}$$

$$\begin{aligned}
\frac{u_{2i}^2 q^{-1} - c_-^2 q}{u_{2i}^{-2} - c_-^2 q^2} (-1)^{M_2+1} &= \prod_{j=1}^{M_1} \frac{(\frac{u_{2i}}{u_{1j}} q^{-\frac{1}{2}} - \frac{u_{1j}}{u_{2i}} q^{\frac{1}{2}})(u_{1j} u_{2i} q^{-\frac{1}{2}} - \frac{q^{\frac{1}{2}}}{u_{1j} u_{2i}})}{(\frac{u_{2i}}{u_{1j}} q^{\frac{1}{2}} - \frac{u_{1j}}{u_{2i}} q^{-\frac{1}{2}})(u_{1j} u_{2i} q^{\frac{1}{2}} - \frac{q^{-\frac{1}{2}}}{u_{1j} u_{2i}})} \prod_{j=1}^{M_3} \frac{(\frac{u_{3j}}{u_{2i}} q^{-\frac{1}{2}} - \frac{u_{2i}}{u_{3j}} q^{\frac{1}{2}})(u_{3j} u_{2i} q^{\frac{1}{2}} - \frac{q^{-\frac{1}{2}}}{u_{3j} u_{2i}})}{(\frac{u_{3j}}{u_{2i}} q^{\frac{1}{2}} - \frac{u_{2i}}{u_{3j}} q^{-\frac{1}{2}})(u_{3j} u_{2i} q^{-\frac{1}{2}} - \frac{q^{\frac{1}{2}}}{u_{3j} u_{2i}})} \\
&\quad i = 1, \dots, M_2 \\
-1 &= \prod_{j=1}^{M_2} \frac{(\frac{u_{3i}}{u_{2j}} q^{\frac{1}{2}} - \frac{u_{2j}}{u_{3i}} q^{-\frac{1}{2}})(u_{2j} u_{3i} q^{\frac{1}{2}} - \frac{q^{-\frac{1}{2}}}{u_{2j} u_{3i}})}{(\frac{u_{3i}}{u_{2j}} q^{-\frac{1}{2}} - \frac{u_{2j}}{u_{3i}} q^{\frac{1}{2}})(u_{2j} u_{3i} q^{-\frac{1}{2}} - \frac{q^{\frac{1}{2}}}{u_{2j} u_{3i}})} \prod_{j \neq i}^{M_3} \frac{(\frac{u_{3i}}{u_{3j}} q^{-1} - \frac{u_{3j}}{u_{3i}} q)(u_{3i} u_{3j} q^{-1} - \frac{q}{u_{3i} u_{3j}})}{(\frac{u_{3j}}{u_{3i}} q^{-1} - \frac{u_{3i}}{u_{3j}} q)(u_{3i} u_{3j} q - \frac{q^{-1}}{u_{3i} u_{3j}})} \\
&\quad i = 1, \dots, M_3
\end{aligned}$$

The transfer matrix eigenvalues are obtained from (6.58), remarking that $\Lambda(u^{(1)}) = \widehat{\Gamma}^{(1)}(u)$:

$$\begin{aligned}
\Lambda(u^{(1\dots m+n-1)}) &= \sum_{k=1}^{m+n} \widetilde{m}_k(u^{(k+1\dots m+n-1)}) \widehat{\Lambda}_k(u^{(k+1\dots m+n-1)}) \times \\
&\quad \times \prod_{j=1}^{M_k} f_k(u^{(k+1\dots m+n-1)}, u_{kj}) \prod_{j=1}^{M_{k-1}} \widetilde{f}_k(u^{(k+1\dots m+n-1)}, u_{k-1j}), \quad (6.62) \\
\widetilde{m}_{m+n}(u) &= (-1)^{[m+n]} m_{m+n}.
\end{aligned}$$

The Bethe equations (6.61) ensure that $\Lambda(u)$ is analytical, in accordance with the analytical Bethe ansatz. The Bethe vectors take the form:

$$\begin{aligned}
\Phi(\{u\}) &= \widehat{B}_{a_1}^{(1)}(u_{11}) \cdots \widehat{B}_{a_{M_1}}^{(1)}(u_{1M_1}) \widehat{F}_{a_1 \dots a_{M_1}}^{(1)}(\{u\}) \Omega, \quad (6.63) \\
&= \widehat{B}_{a_1}^{(1)}(u_{11}) \cdots \widehat{B}_{a_{M_1}^1}^{(1)}(u_{M_1}^{(1)}) \widehat{B}_{a_2^2}^{(2)}(u_{21}) \cdots \widehat{B}_{a_{M_2}^2}^{(2)}(u_{1M_2}) \cdots \widehat{B}_{a_M^{n+m-1}}^{(n+m-1)}(u_{n+m-1,M}) \Omega^{(n+m-1)}.
\end{aligned}$$

We recall the notation $M = \sum_{j=1}^{n+m-1} M_j$, $\Omega^{(k)} = (e_1^{(k-1)})^{\otimes M_{k-1}} \otimes \Omega^{(k-1)}$, $\Omega^{(1)} = \Omega$ and the auxiliary spaces are indicated according to remark 6.4.

7 Bethe vectors

We present here a generalization to open spin chains of the recursion and trace formulas for Bethe vectors, obtained in [35, 36] (see also [16]) for closed spin chains. To our knowledge, this presentation for open spin chain is entirely new.

7.1 Recursion formula for Bethe vectors

From expression (6.63), we can extract a recurrent form for the Bethe vectors,

$$\Phi_M^{n+m}(\{u\}) = \widehat{B}_{a_1^1}^{(1)}(u_{11}) \cdots \widehat{B}_{a_{M_1}^1}^{(1)}(u_{1M_1}) \widehat{\Psi}_{\{u_1\}}^{(1)} \left(\Phi_{M-M_1}^{n+m-1}(\{u_{(>1)}\}) \right), \quad (7.1)$$

$$\widehat{\Psi}_{\{u_1\}}^{(1)} = v^{(2)} \circ (\tau \otimes \pi_{u_{1M_1}}^{(2)} \otimes \cdots \otimes \pi_{u_{11}}^{(2)}) \circ \Delta^{(M_1)}, \quad (7.2)$$

where π_a is the fundamental representation evaluation homomorphism normalized as in (6.51), $v^{(k)}$ is the application of the highest weight vector $e_1^{(k-1)}$:

$$v^{(k)}(X \Omega^{(k-1)}) = X (e_1^{(k-1)})^{\otimes M_{k-1}} \otimes \Omega^{(k-1)} = X \Omega^{(k)}, \quad (7.3)$$

and τ is the morphism

$$\tau : \begin{array}{ccc} \mathfrak{D}_{\mathfrak{m}-1|\mathfrak{n}} & \rightarrow & \mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}/\mathcal{I}_1 \\ d_{ij}(u) & \mapsto & \widehat{d}_{i+1,j+1}(u) \end{array} . \quad (7.4)$$

If we denote by $[\cdot]_{\mathfrak{m}|\mathfrak{n}}$ the grading used in the $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ superalgebra, the mapping τ corresponds to the identification $[j]_{\mathfrak{m}-1|\mathfrak{n}} = [j+1]_{\mathfrak{m}|\mathfrak{n}}$.

Remark that since all the operators in (7.1) apply on the pseudo-vacuum, one can consider the operators built from τ as belonging to $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ instead of $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}/\mathcal{I}_1$. So by induction we build the Bethe vectors from $\mathfrak{D}_{\mathfrak{m}|\mathfrak{n}}$ generators and R -matrices in auxiliary spaces.

7.2 Supertrace formula for Bethe vectors

We can also write the Bethe vector into a supertrace formula and prove the equivalence with the recursion relation discussed above.

Theorem 7.1 *The Bethe vector (7.1) admit a supertrace formulation. We note $A_1; \dots; A_{\mathfrak{m}+\mathfrak{n}-1}$ the ordered sequence of auxiliary spaces $a_1^1, \dots, a_{M_1}^1; a_1^2, \dots, a_{M_2}^2; \dots; a_1^{\mathfrak{m}+\mathfrak{n}-1}, \dots, a_{M_{\mathfrak{m}+\mathfrak{n}-1}}^{\mathfrak{m}+\mathfrak{n}-1}$.*

$$\Phi_M^{\mathfrak{n}+\mathfrak{m}}(\{u\}) = (-1)^{G_1} \text{str}_{A_1 \dots A_{\mathfrak{m}+\mathfrak{n}-1}} \left(\prod_{i=1}^{\mathfrak{m}+\mathfrak{n}-1} \widehat{\mathbb{D}}_{A_i}^{(i)}(\{u_i\}) E_{\mathfrak{n}+\mathfrak{m}, \mathfrak{n}+\mathfrak{m}-1}^{\otimes M_{\mathfrak{n}+\mathfrak{m}-1}} \otimes \dots \otimes E_{21}^{\otimes M_1} \right) \Omega, \quad (7.5)$$

where

$$\widehat{\mathbb{D}}_{A_i}^{(i)}(\{u_i\}) = \prod_{j=1}^{M_i} \overline{\mathcal{R}}_{A_{<i}, a_j^i}^{(i)}(\{u_{i-1}\}, u_{ij}) \widehat{D}_{a_j^i}^{(i)}(u_{ij}) \mathcal{R}_{a_j^i, A_{<i}}^{(i)}(\{u_{i-1}\}, u_{ij}), \quad (7.6)$$

$$G_k = \sum_{i=k}^{\mathfrak{n}+\mathfrak{m}-2} \frac{M_i(M_i+1)}{2} [i], \quad (7.7)$$

$$\overline{\mathcal{R}}_{A_{<i}, a_j^i}^{(i)}(\{u_{i-1}\}, u_{ij}) = \prod_{b < i}^{\rightarrow} \prod_{c=1}^{\overrightarrow{M_b}} \overline{\mathbb{R}}_{a_j^i a_c^b}^{(i,b)}(u_{ij}, u_{bc}^{(b+1 \dots i-1)}), \quad (7.8)$$

$$\mathcal{R}_{a_j^i, A_{<i}}^{(i)}(\{u_{i-1}\}, u_{ij}) = \prod_{b < i}^{\leftarrow} \prod_{c=1}^{\overleftarrow{M_b}} \mathbb{R}_{a_c^b a_j^i}^{(b,i)}(u_{bc}^{(b+1 \dots i-1)}, u_{ij}). \quad (7.9)$$

Proof: Equivalence is proven along the following lines. Starting from expression (7.5), we can extract the M_1 auxiliary spaces corresponding to the first step of the nested Bethe ansatz :

$$\begin{aligned} \Phi_M^{\mathfrak{n}+\mathfrak{m}}(\{u\}) &= (-1)^{\frac{M_1(M_1+1)}{2}} [1] \text{str}_{A_1} \left[\widehat{\mathbb{D}}_{A_1}^{(1)}(\{u_1\}) \times (-1)^{G_2} \times \right. \\ &\quad \left. \text{str}_{A_2 \dots A_{\mathfrak{m}+\mathfrak{n}-1}} \left(\prod_{i=2}^{\mathfrak{m}+\mathfrak{n}-1} \widehat{\mathbb{D}}_{A_i}^{(i)}(\{u_i\}) E_{\mathfrak{n}+\mathfrak{m}, \mathfrak{n}+\mathfrak{m}-1}^{\otimes M_{\mathfrak{n}+\mathfrak{m}-1}} \otimes \dots \otimes E_{32}^{\otimes M_2} \right) \otimes E_{21}^{\otimes M_1} \right] \otimes \Omega. \end{aligned}$$

Using the isomorphism $End(\mathbb{C}^{m+n}) \sim \mathbb{C}^{m+n} \otimes \mathbb{C}^{m+n}$, one can rewrite, for any $A(v)$, the supertrace with an E_{21} matrix as:

$$\begin{aligned} str\left(\widehat{D}^{(1)}(u) A(v) E_{21}\right) &= \sum_{j=1}^{m+n} (e_1^t \otimes e_j^t \otimes \widehat{d}_{1j}^{(1)}(u)) A(v) (e_1 \otimes e_2 \otimes 1), \\ &= (-1)^{[1]+[1][A]} \widehat{B}^{(1)}(u) A(v) (e_2 \otimes 1). \end{aligned} \quad (7.10)$$

Using formula (7.10) for the auxiliary spaces $1, \dots, M_1$, and remarking that the case $j_a = 1$ for $a = 1, \dots, M_1$ does not contribute, we obtain:

$$\begin{aligned} \Phi_M^{n+m}(\{u\}) &= \widehat{B}_{a_1^1}^{(1)}(u_{11}) \cdots \widehat{B}_{a_{M_1}^1}^{(1)}(u_{1M_1}) (-1)^{G_2} \times \\ &\quad str_{A_2 \dots A_{m+n-1}} \left(\prod_{i=2}^{m+n-1} \widehat{\mathbb{D}}_{A_i}^{(i)}(\{u_i\}) E_{n+m, n+m-1}^{\otimes M_{n+m-1}} \otimes \cdots \otimes E_{32}^{\otimes M_2} \right) \Omega^{(2)}. \end{aligned} \quad (7.11)$$

To end the proof, we remark that:

$$(-1)^{G_2} str_{A_2 \dots A_{m+n-1}} \left(\prod_{i=2}^{m+n-1} \widehat{\mathbb{D}}_{A_i}^{(i)}(\{u_i\}) E_{n+m, n+m-1}^{\otimes M_{n+m-1}} \otimes \cdots \otimes E_{32}^{\otimes M_2} \right) \Omega^{(2)} = \widehat{\Psi}_{\{u_1\}}^{(1)} \left(\Phi_{M-M_1}^{n+m-1}(\{u_{(>1)}\}) \right)$$

which allows to recover the form (7.2). ■

Remark 7.1 (Conjecture) *Although theorem 7.1 has been proven only when $K^+(u)$ belongs to a NABA couple, the expression (7.5) does not depend on $K^+(u)$: we conjecture that this expression is valid for any couple of diagonal $K^\pm(u)$ matrices. This conjecture is supported by the fact that analytical Bethe ansatz is known to work for any diagonal boundary matrices.*

7.3 Examples of Bethe vectors

To illustrate the supertrace formula, we present here some explicit examples of Bethe vectors associated to small numbers of excitations.

Bethe vectors of $\mathcal{D}_{m|n}$ with $n+m=2$ and $M_1=M$. We reproduce here the well-known case obtained with algebraic Bethe ansatz (see also section 5).

$$\Phi_M^2(\{u\}) = (-1)^{M[2]} \widehat{d}_{12}^{(1)}(u_{11}) \cdots \widehat{d}_{12}^{(1)}(u_{1M}) \Omega. \quad (7.12)$$

Note that this expression is also valid when $n+m > 2$, setting $M_1 = M$ and $M_k = 0$, $k > 1$.

Bethe vectors of $\mathcal{D}_{m|n}$ with $n+m=3$, $M_1=1$ and $M_2=1$.

$$\begin{aligned} \Phi_{1,1}^3(\{u\}) &= (-1)^{[1]+[2]+[3]} \frac{\mathbf{b}(u_{11}, u_{21})}{\mathbf{a}_2(u_{11}, u_{21})} \widehat{d}_{12}^{(1)}(u_{11}) \widehat{d}_{23}^{(2)}(u_{21}) \Omega \\ &\quad + (-1)^{[1]+[2]+[3]} \frac{\overline{\mathbf{b}}(u_{21}, u_{11})}{\overline{\mathbf{a}}_2(u_{21}, u_{11})} \frac{\mathbf{w}_{32}(u_{11}, u_{21})}{\mathbf{a}_2(u_{11}, u_{21})} \widehat{d}_{13}^{(1)}(u_{11}) \widehat{d}_{22}^{(2)}(u_{21}) \Omega \\ &\quad + (-1)^{[1]} \frac{\overline{\mathbf{w}}_{23}(u_{21}, u_{11})}{\overline{\mathbf{a}}_2(u_{21}, u_{11})} \frac{\mathbf{b}(u_{11}, u_{21})}{\mathbf{a}_2(u_{11}, u_{21})} \widehat{d}_{13}^{(1)}(u_{11}) \widehat{d}_{33}^{(2)}(u_{21}) \Omega. \end{aligned}$$

Again, this expression is also valid when $\mathbf{n} + \mathbf{m} > 3$, setting $M_k = 0$, $k > 2$.

We also computed the Bethe vectors corresponding to $M_1 = M_2 = M_3 = 1$ and $M_k = 0$, $k > 3$. Their expression is rather long, with 11 different terms: we do not write it here explicitly.

8 Conclusion

In this paper, we have proposed a global treatment of the NBA for **universal transfer matrices** of open spins chains with **NABA couple** of boundary matrices. The modification of the nested Bethe ansatz applicable to diagonal boundary matrices that do not form a NABA couple remains to be found. Since the analytical Bethe ansatz can be performed in this case, such a refinement should be possible.

We have computed a **trace formula** for the Bethe vector of the open chain. This formulation could be a starting point for the investigation of the quantized Knizhnik-Zamolodchikov equation following the work [37]. For such a purpose, the coproduct properties of Bethe vectors for open spin chains remain to be studied. Defining a scalar product and computing the norm of these Bethe vectors is also a point of fundamental interest.

From a different point of view, this trace formula and the mapping between the reflection algebras of different size could be the starting point for the construction of a Drinfeld's current realisation [21] for the reflection algebra in the spirit of [38] on the current realisation of the Bethe vector for the periodic case.

The case of open spin chains with general boundary matrices is also a subject of fundamental interest. A deeper understanding of representations of reflection algebras when the K matrix is not diagonal may be of some help. Alternatively, a different approach using another presentation of the reflection algebra could be the clue to go beyond the results obtained so far. Some works have been done for the $\mathbf{m} + \mathbf{n} = 2$ case in [7], but the general treatment for universal transfer matrices remains an open problem. The functional approach developped in [8] for $\mathbf{m} + \mathbf{n} = 2$ also deserves a generalization, both for universal transfer matrices, and for bigger algebras.

A R and M matrices

We remind the general form of the R -matrices we used in the paper [16]. Note we use a more compact form:

$$R_{12}(u, v) = \mathbf{b}(u, v) \mathbb{I} \otimes \mathbb{I} + \sum_{i,j=1}^{\mathbf{m}+\mathbf{n}} \mathbf{w}_{ij}(u, v) E_{ij} \otimes E_{ji}, \quad (\text{A.1})$$

$$\mathbf{w}_{ij}(u, v) = \begin{cases} \mathbf{a}_i(u, v) - \mathbf{b}(u, v) & \text{for } i = j \\ \mathbf{c}_{ij}(u, v) & \text{otherwise} \end{cases},$$

$$\bar{R}_{12}(u, v) = R_{12}(u, \iota(v)) = \bar{\mathbf{b}}(u, v) \mathbb{I} \otimes \mathbb{I} + \sum_{i,j=1}^{\mathbf{m}+\mathbf{n}} \bar{\mathbf{w}}_{ij}(u, v) E_{ij} \otimes E_{ji}. \quad (\text{A.2})$$

The functions involved in these expressions are given by (with the convention $Y(\mathbf{m}) \equiv Y(\mathbf{m}|0)$, $\hat{\mathcal{U}}_q(\mathbf{m}) \equiv \hat{\mathcal{U}}_q(\mathbf{m}|0)$ and $[b] = 0$, $\forall b$, in these two cases):

For $Y(\mathbf{m}|\mathbf{n})$:

$$\mathbf{b}(u, v) = u - v \quad ; \quad \mathbf{a}_a(u, v) = u - v - (-1)^{[a]} \hbar \quad \text{and} \quad \mathbf{w}_{ab}(u, v) = -(-1)^{[b]} \hbar \quad (\text{A.3})$$

$$\bar{\mathbf{b}}(u, v) = u + v \quad ; \quad \bar{\mathbf{a}}_a(u, v) = u + v - (-1)^{[a]} \hbar \quad \text{and} \quad \bar{\mathbf{w}}_{ab}(u, v) = -(-1)^{[b]} \hbar \quad (\text{A.4})$$

For $\widehat{\mathcal{U}}_q(\mathbf{m}|\mathbf{n})$:

$$\mathbf{b}(u, v) = \frac{u}{v} - \frac{v}{u} \quad ; \quad \mathbf{a}_a(u, v) = \frac{u}{v} q^{1-2[a]} - \frac{v}{u} q^{2[a]-1} \quad (\text{A.5})$$

$$\text{and} \quad \mathbf{w}_{ab}(u, v) = (-1)^{[b]} (q - q^{-1}) \left(\frac{u}{v} \right)^{\text{sign}(b-a)}, \quad a \neq b \quad (\text{A.6})$$

$$\bar{\mathbf{b}}(u, v) = uv - \frac{1}{uv} \quad ; \quad \bar{\mathbf{a}}_a(u, v) = uv q^{1-2[a]} - \frac{1}{uv} q^{2[a]-1} \quad (\text{A.7})$$

$$\text{and} \quad \bar{\mathbf{w}}_{ab}(u, v) = (-1)^{[b]} (q - q^{-1}) (uv)^{\text{sign}(b-a)}, \quad a \neq b \quad (\text{A.8})$$

The matrix M is a diagonal matrix:

$$M = \sum_{i=1}^{\mathbf{m}+\mathbf{n}} m_i E_{ii} \quad \text{with} \quad \begin{cases} m_i = 1 & \text{for } Y(\mathbf{m}|\mathbf{n}) \\ m_i = q^{\mathbf{m}-\mathbf{n}-2k+1} q^{-2[k]+4\sum_{i=1}^k [i]} & \text{for } \widehat{\mathcal{U}}_q(\mathbf{m}|\mathbf{n}) \end{cases} \quad (\text{A.9})$$

B Functions appearing in NBA

The functions are constructed from the three functions appearing in the R -matrix, whose explicit form are given in equations (A.3)-(A.8) above:

$$\begin{aligned} \mathbf{f}_i(u, v) &= \frac{\mathbf{a}_i(v, u) \bar{\mathbf{b}}(u^{(i)}, v^{(i)})}{\mathbf{b}(v, u) \bar{\mathbf{b}}(u, v)} \quad , \quad \widetilde{\mathbf{f}}_{i+1}(u, v) = \frac{\mathbf{a}_{i+1}(u, v) \bar{\mathbf{a}}_{i+1}(u, v)}{\mathbf{b}(u, v) \bar{\mathbf{b}}(u, v)} \\ \mathbf{g}_i(u, v) &= \frac{\mathbf{c}_{i-1i}(u, v) \bar{\mathbf{b}}(v^{(i)}, v^{(i)})}{\mathbf{b}(u, v) \bar{\mathbf{b}}(v, v)} \quad , \quad \widetilde{\mathbf{g}}_{i+1}(u, v) = -(-1)^{[i]+[i+1]} \frac{\mathbf{c}_{k+1k}(u, v) \bar{\mathbf{a}}_{k+1}(u, u)}{\mathbf{b}(u, v) \bar{\mathbf{b}}(u, u)} \quad (\text{B.1}) \\ \mathbf{h}_i(u, v) &= -\frac{\bar{\mathbf{c}}_{i-1i}(u^{(i)}, v^{(i)})}{\bar{\mathbf{b}}(u, v)} \quad , \quad \widetilde{\mathbf{h}}_{i+1}(u, v) = \frac{\bar{\mathbf{c}}_{i+1i}(u^{(i)}, v^{(i)}) \bar{\mathbf{a}}_{i+1}(u, u) \bar{\mathbf{b}}(v^{(i)}, v^{(i)})}{\bar{\mathbf{b}}(u, u) \bar{\mathbf{b}}(v, v) \bar{\mathbf{b}}(u, v)} \end{aligned}$$

We also use (presented here for $\mathcal{A}_{\mathbf{m}|\mathbf{n}} = \widehat{\mathcal{U}}_q(\mathbf{m}|\mathbf{n})$; for $\mathcal{A}_{\mathbf{m}|\mathbf{n}} = Y(\mathbf{m}|\mathbf{n})$ one has to set $q = 1$ in the relations below):

$$\psi_i(u) = \frac{\bar{\mathbf{c}}_{i+1i}(u, u)}{\bar{\mathbf{a}}_i(u, u)} \quad (\text{B.2})$$

$$\chi_k(u) = \begin{cases} -q^{2[k]-1} \frac{\bar{\mathbf{b}}(u, u)}{\bar{\mathbf{b}}(u^{(k)}, u^{(k)})} \eta_k(u, c_+) & \text{for } k = 1 \text{ and } a = 1 \\ -q^{2[k]-1} \frac{\bar{\mathbf{b}}(u, u)}{\bar{\mathbf{b}}(u^{(k)}, u^{(k)})} & \text{else} \end{cases} \quad (\text{B.3})$$

$$\eta_k(u, c_+) = \frac{\bar{\mathbf{b}}(\bar{c}_+, u^{(k+1)})}{\bar{\mathbf{b}}(\bar{c}_+, u^{(k+1)})} \quad (\text{B.4})$$

$$\begin{aligned} \mathbf{e}_k(u) \mathbb{I}_b^{(k)} &= \text{tr}_a(M_a^{(k)} \bar{\mathbb{R}}_{ab}^{(k)}(u, u) \mathbb{R}_{ba}^{(k)}(u, u)) \\ &= q^{-k+1} q^{-2n-2 \sum_{i=k}^{m+n} [i] + 4 \sum_{i=1}^{m+n} [i]} (-1)^{[k]} \frac{\bar{\mathbf{b}}(u^{(\overline{k \dots m+n})}, u^{(\overline{k \dots m+n})})}{\bar{\mathbf{a}}_k(u, u)} \mathbb{I}_b^{(k)} \end{aligned} \quad (\text{B.5})$$

$$\tilde{\mathbf{m}}_k(u) = q^{1-2[k]} \frac{\bar{\mathbf{a}}_{k+1}(u, u) \mathbf{e}_{k+1}(u)}{\bar{\mathbf{b}}(u, u)} \quad \text{for } k \neq 1. \quad (\text{B.6})$$

The following useful relations are used in the paper:

$$\mathbf{b}(u, v) \mathbf{a}_i(\bar{u}^{(j)}, \bar{v}^{(j)}) = \mathbf{a}_i(u, v) \mathbf{a}_j(u, v) - \mathbf{w}_{ij}(u, v) \mathbf{w}_{ji}(u, v), \quad i \neq j, \quad (\text{B.7})$$

$$\begin{aligned} \mathbf{b}(u, v) \mathbf{w}_{ij}(\bar{u}^{(k)}, \bar{v}^{(k)}) &= \mathbf{w}_{ij}(u, v) \mathbf{a}_k(u, v) - \mathbf{w}_{ik}(u, v) \mathbf{w}_{kj}(u, v), \quad i > j > k; \\ \mathbf{b}(u^{(\bar{i})}, v^{(\bar{i})}) &= \mathbf{a}_i(u, v), \end{aligned} \quad (\text{B.8})$$

$$\mathbf{a}_j(u, v) - \mathbf{w}_{ij}(u, v) = \mathbf{b}(u, v) q^{sq(j-i)(-1+2[j])}, \quad (\text{B.9})$$

$$q^{2-4[i]} \psi_{i-1}(u^{(i-1)}) = \psi_{i-1}(u^{(i-1, i)}) - \psi_{i-1}(u^{(i-1, i)}) \psi_i(u^{(i)}) \quad (\text{B.10})$$

References

- [1] I. Cherednik, *Factorizing particles on the half-line and roots systems*, Teor. Mat. Fiz. **61** (1984) 55.
- [2] E.K. Sklyanin, *Boundary conditions for integrable quantum systems*, J. Phys. **A21** (1988) 2375.
- [3] L. Faddeev, *How Algebraic Bethe Ansatz works for integrable model*, in *Symétries Quantiques*, Les Houches summerschool proceedings **64**, Eds A. Connes, K. Gawedzki and J. Zinn-Justin, North-Holland 1998 and [hep-th/9605187](#).
- [4] L.A. Takhtajan and L.D. Faddeev, *The quantum method of the inverse problem and the Heisenberg XYZ model*, Russ. Math. Surveys **34** (1979) 11-68.
- [5] P.P. Kulish and E.K. Sklyanin, *Quantum inverse scattering method and the Heisenberg ferromagnet*, Phys. Lett. **A70** (1979) 461-463;
Solution of the Yang-Baxter equation, J. Sov. Math. **19** (1982) 1596-1620.
- [6] P.P. Kulish and N. Yu. Reshetikhin, *Diagonalisation of $gl(\mathbf{n})$ invariant transfer matrices and quatum N -wave system (Lee model)*, J. Phys. **A16** (1983) L591-L596.

- [7] P. Baseilhac and K. Koizumi, *Exact spectrum of the XXZ open spin chain from the q -Onsager algebra representation theory*, JSTAT **0709** (2007) P09006 and [arXiv:hep-th/0703106];
P. Baseilhac, *New results in the XXZ open spin chain*, proceedings of RAQIS07 (Annecy-le-Vieux, France) and [arXiv:0712.0452].
- [8] A. Kuniba, K. Sakai, J. Suzuki, *Continued fraction TBA and functional relations in XXZ model at root of unity*, Nucl. Phys. **B525**[FS] (1998) 597 and [arXiv:math/9803056];
R. Nepomechie, *Solving the open XXZ spin chain with nondiagonal boundary terms at roots of unity*, Nucl. Phys. **B622** (2002) 615 and [arXiv:hep-th/0110116];
R. Nepomechie, *Bethe Ansatz solution of the open XXZ chain with nondiagonal boundary terms*, J. Phys. **A37** (2004) 433 and [arXiv:hep-th/0304092];
R. Murgan, R. Nepomechie, Chi Shi, *Exact solution of the open XXZ chain with general integrable boundary terms at roots of unity*, JSTAT **0608** (2006) P006 and [arXiv:hep-th/0605223];
L. Frappat, R. Nepomechie, E. Ragoucy, *Complete Bethe Ansatz solution of the open spin- s XXZ chain with general integrable boundary terms*, JSTAT **0709** (2007) P09008 and [arXiv:0707.0653].
- [9] Junpeng Cao, Hai-Qing Lin, Kang-jie Shi, Yupeng Wang, *Exact solutions and elementary excitations in the XXZ spin chain with unparallel boundary fields*, Nucl. Phys. **B663** (2003) 487 and [arXiv:cond-mat/0212163];
- [10] W. Galleas and M.J. Martins, *Solution of the $SU(\mathfrak{n})$ vertex model with non-diagonal open boundaries*, Phys. Lett. **A335** (2005) 167 and [arXiv:nlin.SI/0407027].
- [11] W.L. Yang and Y.Z. Zhang, *Exact solution of the $A_{n-1}^{(1)}$ trigonometric vertex model with non-diagonal open boundaries*, JHEP **0501** (2005) 021 and [arXiv:hep-th/0411190].
- [12] N.Yu. Reshetikhin, *The spectrum of the transfer matrices connected with Kac-Moody algebras*, Lett. Math. Phys. **14** (1987) 235.
- [13] D. Arnaudon, N. Crampé, A. Doikou, L. Frappat and É. Ragoucy, *Analytical Bethe Ansatz for closed and open $gl(\mathfrak{n})$ -spin chains in any representation*, JSTAT **02** (2005) P02007 and [arXiv:math-ph/0411021].
- [14] D. Arnaudon, N. Crampé, A. Doikou, L. Frappat and É. Ragoucy, *Spectrum and Bethe ansatz equations for the $U_q(gl(\mathfrak{n}))$ closed and open spin chains in any representation*, Annales H. Poincaré **7** (2006) 1217 and [arXiv:math-ph/0512037].
- [15] É. Ragoucy and G. Satta, *Analytical Bethe ansatz for closed and open $gl(\mathfrak{m}|\mathfrak{n})$ super-spin chains in arbitrary representations*, JHEP **09** (2007) 001 and [arXiv:hep-th/0706.3327].
- [16] S. Belliard and É. Ragoucy, *The nested Bethe ansatz for 'all' closed spin chains*, J. Phys. **A41** (2008) 295202 and arXiv:0804.2822 [math-ph].

- [17] H.J. de Vega and A. González-Ruiz, *Boundary K-matrices for the six vertex and the $n(2n-1)$ A_{n-1} vertex models*, J. Phys. **A26** (1993) L519 and [arXiv:hep-th/9211114];
Exact solution of the $SU_q(n)$ invariant quantum spin chains, Nucl. Phys. **B417** (1994) 553 and [arXiv:hep-th/9309022];
Exact Bethe ansatz solution of A_{n-1} chains with non- $SU_q(\mathfrak{n})$ invariant open boundary conditions, [arXiv:hep-th/9404141].
- [18] R.H. Yue, H. Fan, B. Hou, *Exact diagonalisation of the quantum supersymmetric $SU_q(\mathfrak{n}|\mathfrak{m})$ model*, Nucl. Phys. **B462** (1996) 167 and [arXiv:cond-mat/9603022].
- [19] N. Yu. Reshetikhin, L. A. Takhtajan and L. D. Faddeev, *Quantization of Lie Groups and Lie algebras*, Leningrad Math. J. **1** (1990) 193–225.
- [20] V. G. Drinfeld, *Quantum groups*, in ‘Proc. Int. Congress Math., Berkeley, 1986’, AMS, Providence RI, 1987, pp. 798–820.
- [21] V. G. Drinfeld, *A new realization of Yangians and quantized affine algebras*, Soviet Math. Dokl. **36** (1988) 212–216.
- [22] M. Jimbo, *A q difference analog of $U(G)$ and the Yang-Baxter equation*, Lett. Math. Phys. **10** (1985) 63;
A q -difference analogue of $U(\mathfrak{gl}(n+1))$, Hecke algebra and the Yang-Baxter equation, Lett. Math. Phys. **11** (1986) 247.
- [23] N. Reshetikhin, M. Semenov-Tian-Shansky, *Central extensions of quantum current groups*, Lett. Math. Phys. **19** (1990) 133.
- [24] D. Arnaudon, J. Avan, N. Crampé, A. Doikou, L. Frappat and É. Ragoucy, *General boundary conditions for the $sl(\mathfrak{n})$ and $sl(\mathfrak{m}|\mathfrak{n})$ open spin chains*, JSTAT **0408** (2004) P005 and [arXiv:math-ph/0406021].
- [25] G.M. Gaudenberger, *New non-diagonal solutions to the $A_n^{(1)}$ boundary Yang–Baxter equation*, [arXiv:hep-th/9911178].
- [26] A. Molev and É. Ragoucy, *Representations of reflection algebras*, Rev. Math. Phys. **14** (2002) 317 and [arXiv:math-qa/0406021].
- [27] V. Chari and A. Pressley, *Yangians and R-matrices*, L’Enseignement Math. **36** (1990) 267;
A guide to quantum groups, Cambridge University Press 1994.
- [28] A.I. Molev, *Irreducibility criterion for tensor products of Yangian evaluation modules*, Duke Math. J. **112** (2002) 307 and [arXiv:math.QA/0009183].
- [29] R.B. Zhang, *The $gl(\mathfrak{m}|\mathfrak{n})$ super-Yangian and its finite dimensional representations*, Lett. Math. Phys. **37** (1996) 419 and [arXiv:q-alg/9507029];
Finite dimensional irreducible representations of the supergroup $U_q(\mathfrak{gl}(\mathfrak{m}|\mathfrak{n}))$, J. Math. Phys. **34** (1993) 1236.

- [30] M. Rosso, *Finite dimensional representations of the quantum analogue of the enveloping algebra of a complex simple Lie algebra*, Commun. Math. Phys. **117** (1988) 581.
- [31] A.I. Molev, *Yangians and their applications*, Handbook of Algebra **3** (M. Hazewinkel, Ed.) Elsevier (2003) 907 and [arXiv:math.QA/021128].
- [32] M. Nazarov, *Quantum Berezinian and the classical Capelli identity*, Lett. Math. Phys. **21** (1991) 123.
- [33] S. Belliard and É. Ragoucy, *Berezinian for $\widehat{U}_q(gl(\mathfrak{m}|\mathfrak{n}))$* , in preparation.
- [34] E. Sklyanin, L. Takhtajan and L. Faddeev, *The Quantum Inverse Problem Method*, Theor. Math. Phys. **40** (1979) 688.
- [35] E. Mukhin, V. Tarasov and A. Varchenko, *Bethe eigenvectors of higher transfer matrices*, JSTAT **0608** (2006) P08002 and [arXiv:math.QA/0605015].
- [36] V. Tarasov and A. Varchenko, *Combinatorial formulae for nested Bethe vector*, [arXiv:math.QA/0702277].
- [37] V. Tarasov and A. Varchenko, *Solutions to the quantized Knizhnik-Zamolodchikov equation and the Bethe ansatz*, [arXiv:hep-th/9411181].
- [38] S. Khoroshkin and S. Pakuliak, *Weight function for $U_q(\widehat{\mathfrak{sl}}_3)$* , Theor. Math. Phys. **145** (2005) 1373 ;
A computation of an universal weight function for the quantum affine algebra $U_q(\widehat{gl}_N)$, arXiv:0711.2819 [math.QA].